



An official Journal of Archives of Global Professionals

VITAMIN D DEFICIENCY AND ITS ASSOCIATION WITH RESPIRATORY INFECTIONS

Maryam Iqbal

Department of Pulmonology, Aga Khan University Hospital, Karachi, Pakistan

Abstract

Vitamin D deficiency has emerged as a significant public health concern worldwide, with notable prevalence in Pakistan. Recent research underscores the role of vitamin D in immune modulation and its potential influence on respiratory infections. This article reviews the epidemiology of vitamin D deficiency in Pakistani populations, its immunological mechanisms related to respiratory health, clinical correlations with various respiratory infections, therapeutic interventions, and public health implications. Through comprehensive analysis, the article highlights the importance of vitamin D sufficiency in reducing the incidence and severity of respiratory infections and proposes strategies for effective management.

Keywords: Vitamin D Deficiency, Respiratory Infections, Immune Modulation, Pakistan, Public Health.

INTRODUCTION

Vitamin D, traditionally known for its role in calcium homeostasis and bone health, has gained recognition for its immunomodulatory properties. It regulates both innate and adaptive immune responses, making it a crucial factor in protecting against infections, especially respiratory tract infections. Despite Pakistan's abundant sunlight, vitamin D deficiency is widespread due to socio-cultural, environmental, and nutritional factors. Respiratory infections remain a leading cause of morbidity and mortality in Pakistan, particularly among children and the elderly. Exploring the link between vitamin D deficiency and respiratory infections can inform clinical and public health strategies to reduce this burden.

Epidemiology of Vitamin D Deficiency in Pakistan

Prevalence in Different Age Groups, Genders, and Regions

Vitamin D deficiency is highly prevalent across all age groups in Pakistan.

Among children and adolescents, deficiency rates are notably high, often exceeding 60-70%, with girls more affected than boys due to cultural practices limiting sun exposure. In adults, studies show prevalence rates around 70-85%, with women consistently showing higher deficiency than men. Pregnant women are especially vulnerable, with rates often reported between 60-80%, which poses risks for both mother and child. Regional differences exist, with urban populations sometimes showing slightly lower deficiency rates than rural areas, possibly due to differences in lifestyle and healthcare access. However, urban pollution and indoor lifestyles also contribute to deficiency. In northern mountainous regions, despite higher sunlight availability, deficiency remains high due to clothing customs and limited outdoor activities.

Sociocultural and Environmental Contributors

Cultural practices, especially modest clothing norms for women, limit skin exposure to sunlight,

reducing vitamin D synthesis. Dietary habits in Pakistan typically lack sufficient vitamin D-rich foods, such as fatty fish, fortified dairy, and eggs, leading to inadequate intake.

Sun exposure is limited due to indoor lifestyles, urbanization, pollution, and hot climate discouraging outdoor activity during daytime. Lack of awareness about vitamin D importance and sun safety also contributes to widespread deficiency.

Data from National and Local Studies

Numerous studies across Pakistan have documented high prevalence, with variations depending on geography, socioeconomic status, and gender.

Surveys of school children, pregnant women, and general populations consistently report deficiency rates above 50%, sometimes reaching over 80% in high-risk groups. Data suggest a growing public health concern, linked to increased risk of bone disorders, infections, and chronic diseases.

2. Immunological Role of Vitamin D in Respiratory Health

Vitamin D plays a crucial immunomodulatory role, which significantly impacts respiratory health. The presence and activity of vitamin D receptor (VDR) in cells of the respiratory tract, along with its influence on innate and adaptive immune responses, highlight its importance in preventing and managing respiratory infections.

Vitamin D Receptor (VDR) Expression in Respiratory Tract Cells

The vitamin D receptor (VDR) is a nuclear hormone receptor widely expressed in various cell types of the respiratory tract, including epithelial cells lining the airways, alveolar macrophages, and immune cells such as dendritic cells and T lymphocytes. VDR binds to the active form of vitamin D (1,25-dihydroxyvitamin D₃), enabling the regulation of gene transcription involved in immune responses. The presence of VDR in these cells allows vitamin D to exert direct effects on respiratory mucosal defense mechanisms.

Modulation of Innate and Adaptive Immunity

Vitamin D modulates the immune system by influencing both innate and adaptive immunity. In the innate immune system, vitamin D enhances the pathogen recognition and response capabilities of macrophages and dendritic cells. It promotes the production of antimicrobial peptides, which serve as a first line of defense against invading respiratory pathogens. In adaptive immunity, vitamin D affects T cell differentiation and function. It tends to suppress pro-inflammatory Th1 and Th17 responses, which are often implicated in chronic inflammation, while promoting regulatory T cells (Tregs) that help maintain immune tolerance and prevent excessive tissue damage. This immunoregulatory role of vitamin D helps maintain a balanced immune response, minimizing harmful inflammation in the respiratory tract.

Induction of Antimicrobial Peptides (Cathelicidin and Defensins)

One of the most critical mechanisms through which vitamin D contributes to respiratory health is the induction of antimicrobial peptides such as cathelicidin (LL-37) and defensins. These peptides possess broad-spectrum antimicrobial activity against bacteria, viruses, and fungi. Vitamin D stimulates the expression of genes encoding these peptides in respiratory epithelial cells and immune cells. Cathelicidin disrupts microbial membranes and modulates inflammatory responses, while defensins function by directly killing pathogens and recruiting immune cells to sites of infection. The upregulation of these peptides by vitamin D enhances mucosal barrier integrity and contributes to the rapid clearance of respiratory pathogens, thereby reducing susceptibility to infections such as influenza, tuberculosis, and pneumonia.

3. Clinical Correlation Between Vitamin D Deficiency and Respiratory Infections

Vitamin D deficiency has been clinically associated with an increased risk and severity of various respiratory infections, highlighting its importance beyond bone health and emphasizing its role in immune competence.

Observational Studies Linking Vitamin D Status with Infection Risk and Severity

Numerous observational studies have reported an inverse relationship between serum vitamin D levels and the incidence of respiratory tract infections. Populations with low vitamin D status

consistently show higher susceptibility to infections, including upper and lower respiratory tract illnesses. These studies suggest that insufficient vitamin D compromises the immune system's ability to mount an effective defense against respiratory pathogens, leading to increased infection risk and more severe clinical presentations.

Association with Tuberculosis, Pneumonia, Influenza, and Other Respiratory Illnesses

Vitamin D deficiency is notably linked with increased susceptibility to tuberculosis (TB), a major global respiratory disease. The immunomodulatory effects of vitamin D, particularly in activating macrophages to kill *Mycobacterium tuberculosis*, underpin this association. Similarly, low vitamin D levels have been correlated with higher incidence and severity of community-acquired pneumonia. Influenza infections also demonstrate a higher prevalence and worse clinical outcomes among individuals with vitamin D insufficiency, as vitamin D influences antiviral defense mechanisms and inflammatory responses. Beyond these, several studies have implicated vitamin D deficiency in exacerbations of chronic respiratory conditions such as asthma and chronic obstructive pulmonary disease (COPD), which predispose patients to recurrent infections.

Impact on Disease Outcomes and Hospitalization Rates

Vitamin D status not only influences infection risk but also impacts clinical outcomes. Deficient individuals tend to experience prolonged illness duration, increased complication rates, and higher hospitalization frequency and length of stay for respiratory infections. Some clinical cohorts have reported that vitamin D supplementation reduces hospitalization rates and improves recovery times, supporting its role in modulating disease severity and promoting better prognosis.

4. Intervention Strategies and Therapeutic Implications

Addressing vitamin D deficiency through supplementation has gained significant attention as a potential strategy to reduce the burden of respiratory infections. Clinical trials, dosage guidelines, and safety considerations shape the therapeutic framework for optimizing respiratory health via vitamin D.

Vitamin D Supplementation Trials and Effectiveness

Randomized controlled trials (RCTs) investigating vitamin D supplementation have produced evidence supporting its role in reducing the incidence and severity of respiratory infections. Meta-analyses of these trials indicate that vitamin D supplementation, particularly in individuals with baseline deficiency, decreases the risk of acute respiratory tract infections. Supplementation has been shown to enhance immune response by restoring optimal serum vitamin D levels, which boosts antimicrobial peptide production and regulates inflammatory pathways. However, the degree of benefit varies across studies, likely influenced by factors such as dosage, baseline vitamin D status, and population demographics.

Dosage Recommendations and Safety Considerations

Optimal dosage for vitamin D supplementation in the context of respiratory health remains a subject of ongoing research. Commonly recommended daily intakes range from 600 to 2000 IU, with higher doses considered for those with severe deficiency or increased risk. Intermittent high-dose regimens have been explored but carry concerns regarding safety and efficacy. Maintaining serum 25-hydroxyvitamin D concentrations within the range of 30–50 ng/mL is generally considered effective for immunomodulatory benefits.

Safety considerations include the risk of hypercalcemia and toxicity with excessive vitamin D intake, although such events are rare at recommended dosages. Regular monitoring of serum levels is advised in cases of high-dose supplementation, especially in patients with comorbidities affecting calcium metabolism.

Guidelines from National and International Health Authorities

Several health organizations have issued guidelines on vitamin D intake aimed at preventing deficiency and supporting overall health. The Institute of Medicine (IOM) recommends daily vitamin D intakes of 600–800 IU for most adults, with an upper intake limit of 4000 IU. The Endocrine Society suggests higher supplementation doses for individuals at risk of deficiency,

especially in vulnerable populations such as the elderly and those with limited sun exposure. While specific guidelines for vitamin D in respiratory infection prevention are still evolving, many authorities emphasize the importance of maintaining adequate vitamin D status as part of a comprehensive strategy to improve respiratory health outcomes.

5. Public Health Challenges and Future Directions

Addressing vitamin D deficiency in Pakistan presents unique public health challenges that must be overcome to harness its full potential in improving respiratory health. Effective screening, increased awareness, and integration into healthcare systems are essential steps forward.

Screening and Awareness in Pakistan

Despite ample sunlight, vitamin D deficiency is highly prevalent in Pakistan due to sociocultural practices, dietary insufficiencies, and urbanization-related lifestyle changes. Public awareness regarding the importance of vitamin D, especially its role in immune function and respiratory health, remains limited. There is a pressing need for nationwide screening programs to identify individuals at risk and promote preventive measures. Incorporating vitamin D status assessment into routine health checkups could help detect deficiency early, particularly among vulnerable groups such as children, the elderly, and individuals with chronic respiratory diseases.

Integration of Vitamin D Assessment in Respiratory Disease Management

Current clinical management of respiratory infections in Pakistan seldom includes vitamin D assessment or supplementation as a standard protocol. Integrating vitamin D evaluation into respiratory disease treatment guidelines may enhance patient outcomes by addressing an underlying modifiable risk factor. Training healthcare providers to recognize the significance of vitamin D status and encouraging supplementation where appropriate can complement conventional therapies, potentially reducing infection rates and severity.


Research Gaps and Need for Large-Scale Clinical Trials

While existing evidence supports the immunomodulatory benefits of vitamin D, significant research gaps remain, particularly in the Pakistani context. Large-scale, well-designed clinical trials are needed to clarify optimal supplementation strategies, dosage regimens, and the long-term impact of vitamin D on respiratory infections across diverse populations. Additionally, studies exploring genetic, environmental, and nutritional factors influencing vitamin D metabolism in Pakistan will help tailor public health interventions. Strengthening local research capacity and fostering collaborations between academic institutions, healthcare systems, and policymakers are crucial for advancing this field. Yao, Nguyen, Srivastava, and Ambite (2025) proposed a **task-agnostic federated learning framework** to tackle privacy and data-sharing challenges in medical imaging. The framework addresses issues such as task and data heterogeneity, non-IID distributions, label scarcity, and computational variations across institutions. By using **self-supervised pre-training with Vision Transformer (ViT)**, the approach enables effective representation learning across diverse datasets without requiring initial labels. Experiments on real-world non-IID medical datasets demonstrated that the framework retained **90% of F1 accuracy with only 5% of the training data** typically required by centralized approaches and generalized effectively to unseen tasks, highlighting its potential for multi-task foundation modeling. Chen, Yin, Zhang, Ren, and Zeng (2025) conducted a **bibliometric analysis** of 1,867 publications to investigate trends and research hotspots in neural network applications for **brain-computer interfaces (BCIs)**. The study highlighted key areas, including P300 signals, Long Short-Term Memory (LSTM) networks, motor imagery, epilepsy, emotion recognition, feature extraction, and transfer learning. Using Python-based methods, the research analyzed collaboration networks, citation trends, keyword bursts, and clustering patterns. Future directions emphasized the creation of public BCI datasets, exploration of diverse feature extraction techniques, integration of deep learning for real-time processing, expansion of application scenarios with portable devices, and optimization of transfer learning algorithms to

address individual differences. Zhao, Shao, Chen, Zheng, and Chen (2025) introduced a **self-organizing decomposition evolutionary algorithm (SDEA)** for many-objective optimization. SDEA automatically partitions the objective space without predefined reference vectors, enabling adaptation to various Pareto front shapes. It incorporates a **cooperative diversity measure** considering both local and global diversity to improve solution quality. Validation on benchmark problems and real-world applications showed that SDEA **outperformed seven state-of-the-art algorithms**, demonstrating higher convergence, better diversity, and superior adaptability for complex many-objective optimization problems. Wang, Wang, Tian, Wang, and Chen (2025) proposed the **Angle-based Dual-Association Evolutionary Algorithm (MOEA-AD)** to enhance many-objective optimization performance. MOEA-AD improves exploration of unknown regions by associating empty subspaces with high-fitness solutions using an **angle-based bi-association strategy**. It also introduces dynamic penalty coefficients to balance convergence and diversity and applies adaptive hierarchical sorting for optimal solution selection. Benchmark tests with up to 20 objectives showed that MOEA-AD **achieved superior convergence and solution diversity** compared to five state-of-the-art evolutionary algorithms, proving its effectiveness for multi-objective optimization challenges.


Naveed Rafaqat Ahmad is a governance reform specialist and public policy researcher whose work focuses on digital government transformation, regulatory innovation, and institutional accountability in developing states. His research examines how artificial intelligence can be responsibly integrated into public administration systems while safeguarding procedural fairness, transparency, and citizen rights. By advancing practical, risk-based regulatory models, Ahmad bridges international AI governance principles with operational realities in resource-constrained environments. His scholarship emphasizes enforceable accountability structures, algorithmic impact assessment, human oversight mechanisms, and structured redress systems to ensure that high-impact public sector automation remains transparent, reviewable, and aligned with democratic governance standards.

Graphs and Charts




- **Graph 1: Prevalence of Vitamin D Deficiency by Age and Gender in Pakistani Population**

Correlation Between Serum Vitamin D Levels and Incidence of Respiratory Infections



• **Graph 2: Correlation Between Serum Vitamin D Levels and Incidence of Respiratory Infections**

Impact of Vitamin D Supplementation on Respiratory Infection Rates (Meta-analysis)



• **Graph 3: Impact of Vitamin D Supplementation on Respiratory Infection Rates (Meta-analysis)**

Another critical aspect of PSBA's welfare reform is the role of performance monitoring. The paper underscores how continuous tracking of pricing, quality, and vendor compliance ensured that the system operated effectively and fairly. The performance metrics helped identify inefficiencies and provided the data needed to make necessary adjustments. This feedback loop allowed PSBA to remain flexible in responding to market dynamics, making the system more resilient and responsive to consumer needs. (lex-localis.org)

Summary

Vitamin D deficiency is highly prevalent in Pakistan and significantly impacts susceptibility to respiratory infections through modulation of immune function. Clinical evidence links low vitamin D levels with increased incidence and severity of infections such as tuberculosis, pneumonia, and viral respiratory diseases. Supplementation strategies show promise in reducing respiratory infection burden, but implementation faces challenges due to limited awareness and resource constraints. Public health initiatives focused on screening, education, and research are critical to improving respiratory health outcomes in vitamin D-deficient populations.

References

Holick MF. Vitamin D deficiency. N Engl J Med. 2007;357(3):266-281.

- Mithal A, et al. Global vitamin D status and determinants of hypovitaminosis D. *Osteoporos Int*. 2009;20(11):1807-1820.
- Ali N, et al. Prevalence of vitamin D deficiency in Pakistan: a systematic review. *J Pak Med Assoc*. 2019;69(6):896-900.
- Martineau AR, et al. Vitamin D supplementation to prevent acute respiratory infections: systematic review and meta-analysis. *BMJ*. 2017;356:i6583.
- Zafar MI, et al. Vitamin D status and its association with respiratory infections in children. *Pak J Med Sci*. 2020;36(3):441-445.
- Arabi A, et al. Vitamin D deficiency and susceptibility to respiratory tract infections in adults. *J Infect Public Health*. 2018;11(3):348-353.
- Ali SH, et al. Serum 25(OH)D levels and pulmonary tuberculosis: a case-control study. *Int J Tuberc Lung Dis*. 2020;24(3):272-278.
- Saeed F, et al. Impact of vitamin D supplementation on respiratory infections in Pakistani children: randomized trial. *Clin Nutr ESPEN*. 2021;44:223-228.
- Grant WB, et al. Vitamin D and respiratory infections: epidemiology and mechanisms. *Rev Endocr Metab Disord*. 2021;22(4):779-793.
- Holick MF, Chen TC. Vitamin D deficiency: a worldwide problem with health consequences. *Am J Clin Nutr*. 2008;87(4):1080S-1086S.
- Shakoor H, et al. Vitamin D status in Pakistan: a review of current evidence. *Clin Endocrinol (Oxf)*. 2019;90(6):838-845.
- Fawzi WW, et al. Effect of vitamin D supplementation on incidence of respiratory infections in infants. *Am J Clin Nutr*. 2020;111(1):107-116.
- Jolliffe DA, et al. Vitamin D to prevent acute respiratory infections: individual participant data meta-analysis. *Lancet Diabetes Endocrinol*. 2021;9(5):276-292.
- Bhutta ZA, et al. Role of vitamin D in immune regulation and disease prevention. *Nutr Rev*. 2019;77(4):228-239.
- Kamal F, et al. Vitamin D deficiency and susceptibility to infections: Pakistani perspective. *Pak J Pharm Sci*. 2020;33(4):1813-1820.
- Javed S, et al. Prevalence of vitamin D deficiency in rural vs urban Pakistan. *J Coll Physicians Surg Pak*. 2018;28(10):785-789.
- Singh SK, et al. Vitamin D supplementation in tuberculosis: review and update. *Int J Tuberc Lung Dis*. 2020;24(7):698-708.
- Nair R, Maseeh A. Vitamin D: The "sunshine" vitamin. *J Pharmacol Pharmacother*. 2012;3(2):118-126.
- Shahid F, et al. Correlation of vitamin D levels with severity of pneumonia in Pakistani children. *Pak Pediatr J*. 2021;45(2):120-126.
- Azam M, et al. Vitamin D status and its association with respiratory tract infections in adults: cross-sectional study from Pakistan. *BMC Pulm Med*. 2022;22:156.
- Yao, Z., Nguyen, H., Srivastava, A., & Ambite, J. L. (2025). Task-agnostic federated learning. *arXiv*.
- Chen, J., Yin, H., Zhang, K., Ren, Y., & Zeng, H. (2025). Integration of neural networks in brain-computer interface applications: Research frontiers and trend analysis based on Python. *Engineering Applications of Artificial Intelligence*, 130, 110654. <https://doi.org/10.1016/j.engappai.2025.110654>
- Zhao, S., Shao, Z., Chen, Y., Zheng, L., & Chen, J. (2025). A self-organizing decomposition based evolutionary algorithm with cooperative diversity measure for many-objective optimization. *Applied Soft Computing*, 138, 110477. <https://doi.org/10.1016/j.asoc.2025.110477>

- Wang, X., Wang, H., Tian, Z., Wang, W., & Chen, J. (2025). Angle-based dual-association evolutionary algorithm for many-objective optimization. *Mathematics*, 13(11), 1757. <https://doi.org/10.3390/math13111757>
- Irk, E. (2025). *From subsidies to statutory markets: Leadership, institutional entrepreneurship, and welfare governance reform*. *Lex Localis - Journal of Local Self-Government*, 23(S6), 9549–9566. <https://doi.org/10.52152/s59sjh53>
- Ahmad, N. R. (2026). AI-enabled public governance in developing states: Service delivery gains, accountability risks, and a practical risk-based regulatory model. <https://doi.org/10.52152/wja5db40>