



Multi Omics Profiling of Macrophage Polarization in Ischemic Tissue Repair Repair Mechanisms

Stella Bennett, Victor Parker, Rina Kawamura, Maya Fleming

Mathematical, Physical and Life Sciences Division, University of Oxford, Oxford, United Kingdom

Abstract

Ischemic tissue injury represents a leading cause of global morbidity and mortality, necessitating advanced therapeutic interventions aimed at promoting functional recovery. A critical determinant of successful tissue repair is the dynamic regulation of macrophage polarization, a process wherein macrophages transition from an initial pro-inflammatory phenotype to a subsequent pro-resolving and regenerative state. Traditional studies have predominantly relied on single-layer biological analyses, which often fail to capture the complex, interconnected regulatory networks governing this phenotypic plasticity. This paper presents a comprehensive examination of macrophage polarization in ischemic tissue repair through a multi-omics lens, integrating transcriptomics, epigenomics, proteomics, and metabolomics. By synthesizing data across these molecular strata, we elucidate the profound metabolic reprogramming events, particularly the shifts between glycolysis and oxidative phosphorylation, and their direct mechanistic links to epigenetic landscape remodeling. The analysis underscores how these multi-dimensional datasets can identify novel regulatory hubs and potential therapeutic targets that are entirely obscured in single-omics paradigms. Furthermore, we discuss the methodological challenges and computational strategies required to harmonize heterogeneous datasets. Ultimately, the integration of systems biology into macrophage immunology provides a robust framework for designing next-generation immunomodulatory therapies for ischemic cardiovascular and peripheral vascular diseases.

Keywords: *Ischemia, Macrophage Polarization, Multi-Omics, Systems Biology*

1. Introduction

1.1 The Burden of Ischemic Diseases

Ischemic diseases, encompassing conditions such as myocardial infarction, ischemic stroke, and critical limb ischemia, constitute a profound burden on healthcare systems globally. These pathologies share a common etiological foundation characterized by the acute or chronic restriction of blood supply to tissues, leading to severe hypoxia, nutrient deprivation, and subsequent cellular necrosis [1]. The initial ischemic insult triggers a cascade of detrimental events, including the robust generation of reactive oxygen species, profound intracellular calcium overload, and the catastrophic collapse of mitochondrial bioenergetics [2]. In the immediate aftermath of an ischemic event, the host immune system mounts a highly orchestrated sterile inflammatory response aimed at clearing necrotic debris and initiating the preliminary stages of wound healing. However, the exact trajectory of this inflammatory response dictates the ultimate fate of the affected tissue [3]. An excessive or prolonged inflammatory phase inevitably exacerbates tissue damage, promoting adverse remodeling, fibrosis, and organ dysfunction. Conversely, a tightly regulated and properly timed resolution of inflammation fosters angiogenesis, extracellular matrix reorganization, and parenchymal regeneration [4]. Understanding the delicate balance between tissue destruction and repair following ischemia remains a paramount objective in

contemporary cardiovascular and regenerative medicine. Despite decades of intensive research, the clinical translation of therapies aimed at modulating post-ischemic inflammation has met with limited success, largely due to a fragmented understanding of the cellular and molecular networks that govern the immune response in the ischemic microenvironment [5].

1.2 Macrophage Polarization in Tissue Repair

At the center of the post-ischemic immune response is the macrophage, a highly plastic myeloid lineage cell that plays an indispensable role in orchestrating both the initiation and the resolution of inflammation [6]. Following an ischemic insult, circulating monocytes rapidly infiltrate the damaged tissue in response to damage-associated molecular patterns and local chemokines, subsequently differentiating into tissue macrophages. Historically, the functional diversity of these cells has been conceptualized through a binary classification system, dividing them into classical pro-inflammatory or alternative pro-resolving phenotypes [7]. The early-arriving macrophages typically exhibit the classical phenotype, characterized by the robust secretion of pro-inflammatory cytokines, the production of reactive nitrogen species, and specialized functions directed toward the phagocytosis of necrotic cellular debris [8]. While this early inflammatory phase is essential for clearing the damaged area, its persistence is highly deleterious. Within a few days post-injury, a critical transition must occur wherein the macrophage population shifts toward the alternative phenotype [9]. These pro-resolving macrophages dampen inflammation through the secretion of anti-inflammatory cytokines, secrete vascular endothelial growth factor to promote the formation of new blood vessels, and release transforming growth factor to stimulate fibroblast activity and tissue structural repair [10]. The failure to execute this phenotypic transition is a hallmark of chronic, non-healing ischemic wounds and adverse cardiac remodeling. However, recent advances in single-cell technologies have revealed that this binary paradigm is an immense oversimplification. In reality, tissue macrophages exist across a highly diverse, continuous spectrum of activation states, simultaneously expressing markers of both inflammation and repair depending on spatial and temporal microenvironmental cues [11]. The regulatory mechanisms that dictate a macrophage position along this continuum are extraordinarily complex, integrating signals from the extracellular matrix, local oxygen tensions, and the accumulation of specific metabolites.

1.3 The Need for Multi-Omics Approaches

To fully comprehend the extraordinary plasticity of macrophages in the ischemic environment, it has become increasingly evident that traditional reductionist approaches, which focus on single genes or isolated molecular pathways, are fundamentally inadequate [12]. The transition from an inflammatory to a regenerative macrophage state is not dictated by a single master regulator but is rather the emergent property of thousands of interacting molecules spanning multiple biological layers. Early systems biology approaches typically focused on transcriptomics, utilizing bulk RNA sequencing to identify differential gene expression profiles between early and late post-ischemic time points [13]. While these studies provided valuable compendiums of gene expression, they failed to account for post-transcriptional modifications, protein degradation rates, and the metabolic fluxes that actually execute cellular function [14]. Furthermore, epigenetic modifications, which dictate chromatin accessibility and prime cells for rapid transcriptional responses, act as critical gatekeepers of macrophage fate, yet they are completely invisible to standard transcriptomic assays [15]. Consequently, the contemporary frontier of macrophage immunology demands the implementation of multi-omics profiling. By simultaneously capturing transcriptomic, epigenomic, proteomic, and metabolomic data from the exact same biological contexts, researchers can construct comprehensive, multi-dimensional networks. This integrative approach allows for the identification of causal chains of events, for instance, how the accumulation of a specific local metabolite might alter the activity of chromatin-modifying enzymes, thereby reshaping the epigenetic landscape and ultimately reprogramming the functional proteome of the macrophage.

2. Literature Review and Background

2.1 Pathophysiology of Ischemic Tissue Injury

The sequence of biological events following ischemic injury is characterized by a high degree of spatial

and temporal complexity. When arterial occlusion abruptly halts the delivery of oxygen and metabolic substrates, the affected parenchymal cells rapidly deplete their intracellular reserves of adenosine triphosphate [16]. This catastrophic failure of energy metabolism disables critical transmembrane ion pumps, resulting in cellular swelling, organelle dysfunction, and ultimately necrotic cell death. The rupturing of these necrotic cells releases an immense quantity of intracellular constituents into the extracellular space. These molecules, which include nuclear DNA, high mobility group box proteins, and heat shock proteins, act as potent damage-associated molecular patterns [17]. These patterns are rapidly detected by pattern recognition receptors, particularly toll-like receptors, residing on the surface of local tissue-resident macrophages and endothelial cells. The activation of these receptors initiates a massive signaling cascade, leading to the rapid nuclear translocation of transcription factors that drive the expression of potent chemokines and adhesion molecules [18]. This creates a powerful chemotactic gradient that recruits massive numbers of innate immune cells, primarily neutrophils and inflammatory monocytes, from the bone marrow and spleen to the site of injury. The microenvironment during this acute phase is highly hostile, characterized by severe hypoxia, a highly acidic potential of hydrogen, and an overwhelming presence of destructive proteases and reactive oxygen species. Understanding how newly recruited macrophages survive and adapt to this extreme environment is a critical area of investigation, as their early functional orientation sets the trajectory for the entire subsequent repair process [19].

2.2 Dynamics of Macrophage Phenotypes

The temporal dynamics of macrophage infiltration and phenotypic adaptation are tightly choreographed during successful tissue repair. In murine models of myocardial infarction and hindlimb ischemia, the peak infiltration of inflammatory monocytes occurs within the first three days post-injury [20]. These monocytes rely heavily on specific chemokine receptor axes for their recruitment and rapidly differentiate into macrophages that exhibit high phagocytic capacity and robust expression of inflammatory mediators. Interestingly, these cells also upregulate survival pathways that allow them to function in the nearly anoxic conditions of the infarct core [21]. As the initial necrotic debris is cleared, the local microenvironment begins to change. The precise triggers for macrophage phenotypic transition are multifaceted, involving the phagocytosis of apoptotic neutrophils, the accumulation of specific lipid mediators, and the presence of locally generated anti-inflammatory cytokines [22]. By day seven post-injury, the dominant macrophage population typically exhibits a regenerative phenotype. These cells are essential for coordinating the formation of granulation tissue. They stimulate endothelial cells to sprout and form new capillary networks, a process critical for restoring oxygen delivery to the healing tissue [23]. Furthermore, they communicate extensively with local fibroblasts, regulating the deposition of collagen and other extracellular matrix components to form a stable structural scar. If this transition is delayed or prevented by underlying comorbidities, such as diabetes or chronic hyperlipidemia, the tissue remains locked in a state of chronic inflammation, leading to excessive tissue destruction, pathological fibrosis, and ultimately organ failure [24].

2.3 Single-Omics versus Multi-Omics Paradigms

Historically, the characterization of these diverse macrophage populations relied on flow cytometry and single-omics techniques, predominantly bulk transcriptomics. While transcriptomics revolutionized the field by enabling the simultaneous measurement of thousands of messenger RNA transcripts, it inherently assumes that messenger RNA abundance is a reliable proxy for protein expression and cellular function [25]. Numerous studies have now definitively shown that in highly dynamic environments like the ischemic wound, the correlation between transcript levels and actual protein abundance is surprisingly weak. This discrepancy is driven by complex post-transcriptional regulatory mechanisms, including the action of micro RNAs, dynamic regulation of messenger RNA translation rates, and variable protein degradation kinetics [26]. Epigenomics, captured through assays such as the assay for transposase-accessible chromatin using sequencing, provides a different but equally incomplete picture, revealing the structural availability of the genome but not confirming whether transcription is actually occurring [27]. Proteomics and metabolomics offer the closest representation of the actual phenotypic state of the cell, but when used in isolation, they provide limited

insight into the upstream regulatory networks that produced those functional states [28]. Multi-omics integration overcomes these fundamental limitations. By vertically integrating data from the genome structure up through to the metabolic output, researchers can map the entire flow of biological information. This paradigm shift allows for the identification of master regulatory nodes that coordinate complex phenotypic changes. For example, a single-omics study might identify a change in a metabolic pathway, but a multi-omics study can trace that metabolic shift back to a specific set of transcription factors whose access to the DNA was altered by an upstream epigenetic modification [29].

2.4 Current Challenges in Systems Biology of Macrophages

Despite the immense theoretical power of multi-omics approaches, the practical implementation in the context of ischemic tissue repair presents significant technical and analytical hurdles. The primary challenge lies in the immense heterogeneity of the ischemic microenvironment [30]. When isolating macrophages from an ischemic organ, the resulting sample represents a mixture of cells residing in the necrotic core, the border zone, and the remote uninjured tissue. Because bulk multi-omics assays average the signals across all these cells, critical spatially restricted phenotypes are frequently obscured [31]. Furthermore, acquiring sufficient high-quality biological material for multiple distinct omics assays from the same rare subpopulation of primary macrophages is extremely difficult, often requiring the pooling of tissue from dozens of animals [32]. Computationally, the integration of multi-omics datasets is an enormously complex task. Transcriptomic datasets typically consist of tens of thousands of features, whereas metabolomic datasets might only confidently identify a few hundred metabolites. These vast differences in data scale, coupled with differing dynamic ranges and technical noise profiles, make traditional statistical correlations highly susceptible to false positives [33]. Developing robust bioinformatics frameworks that can accurately perform dimensionality reduction, identify latent factors spanning multiple data layers, and infer true biological causality remains the most significant bottleneck in the field of systems immunology.

3. Methodology and Analytical Framework

3.1 Experimental Design and Ischemic Models

To investigate the multi-omics profile of macrophage polarization, a highly standardized and rigorous experimental design is essential. The investigations detailed in this methodological framework utilize well-established murine models of ischemic injury, specifically the permanent ligation of the left anterior descending coronary artery to induce myocardial infarction, and the excision of the femoral artery to induce critical hindlimb ischemia. These models are chosen due to their high reproducibility and their strong pathophysiological similarities to human ischemic diseases. Wild-type male mice, standardized for age and genetic background, are subjected to the surgical procedures under rigorous sterile conditions with appropriate anesthetic and analgesic protocols. To capture the dynamic temporal shifts in macrophage polarization, tissue harvesting must be performed at multiple, carefully selected time points. These time points typically include day one and day three to capture the peak inflammatory phase, day seven to capture the transition to the reparative phase, and day fourteen to observe the late stages of tissue remodeling and scar stabilization. Following tissue extraction, the ischemic tissue is rapidly processed to minimize ex vivo artifactual changes in gene expression and metabolism. The tissue is subjected to precise enzymatic digestion using carefully optimized cocktails of collagenases and nucleases to release the cellular components from the extracellular matrix while preserving delicate surface antigens. The resulting single-cell suspensions are then passed through sophisticated fluorescence-activated cell sorting systems. Macrophages are identified and isolated based on the expression of established surface markers, ensuring high purity of the final cellular populations for subsequent multi-omics extractions.

3.2 Transcriptomics and Epigenomics Profiling

Following the isolation of high-purity macrophage populations from the ischemic tissues, the cells are immediately partitioned for the parallel extraction of nucleic acids. For transcriptomic profiling, total RNA is extracted using optimized column-based purification methods that ensure high yield and exceptional RNA integrity. The extracted RNA undergoes rigorous quality control using automated electrophoresis systems. High-quality RNA is then utilized for the construction of sequencing libraries.

To capture a comprehensive view of the transcriptome, including both coding and non-coding RNA species, ribosomal RNA depletion strategies are typically employed rather than standard poly-A selection. The resulting libraries are sequenced on high-throughput platforms to generate massive datasets of short reads. In parallel, a separate aliquot of the sorted macrophages is utilized for epigenomic profiling to assess chromatin accessibility. The isolated nuclei are subjected to an optimized assay for transposase-accessible chromatin. In this procedure, a hyperactive transposase enzyme is utilized to simultaneously fragment and tag the open, transcriptionally accessible regions of the chromatin with sequencing adapters. This technique requires significantly fewer cells than traditional chromatin immunoprecipitation methods, making it highly suitable for the rare macrophage subpopulations isolated from in vivo ischemic models. The resulting libraries are sequenced, and the data provides a high-resolution map of the regulatory landscape, identifying active promoters and distant enhancer elements that govern the transcriptional program of the different macrophage activation states.

Table 1: Key Multi-Omics Modalities and Analytical Techniques for Macrophage Profiling

Omics Layer	Primary Biological Target	Common Analytical Platform	Key Output Metric
Transcriptomics	Total Messenger RNA	High-Throughput RNA Sequencing	Read Counts per Gene
Epigenomics	Chromatin Accessibility	Transposase-Accessible Chromatin Sequencing	Genomic Peak enrichment
Proteomics	Global Protein Abundance	Liquid Chromatography Mass Spectrometry	Peptide Spectral Matches
Metabolomics	Small Intracellular Molecules	Targeted Mass Spectrometry	Metabolite Peak Area

3.3 Proteomics and Metabolomics Integration

To move beyond the predictive layers of nucleic acids and assess the actual functional machinery of the macrophages, proteomics and metabolomics must be seamlessly integrated into the analytical pipeline. For global proteomic analysis, the sorted macrophages undergo robust cellular lysis and protein extraction using highly denaturing buffers to ensure the complete solubilization of all protein classes, including hydrophobic membrane proteins. The extracted proteins are then reduced, alkylated, and enzymatically digested into peptides using highly specific proteases such as trypsin. To enable precise relative quantification across the multiple time points and biological replicates, the peptides are typically labeled using tandem mass tags. These isobaric chemical tags allow for the multiplexing of several samples into a single mass spectrometry run, thereby eliminating run-to-run technical variance. The labeled peptides are fractionated and analyzed using ultra-high-performance liquid chromatography coupled to high-resolution orbitrap mass spectrometers. Simultaneous metabolomic profiling requires rapid quenching of cellular metabolism during the sorting process to prevent the artefactual alteration of highly labile metabolites. The intracellular metabolites are extracted using carefully optimized mixtures of cold organic solvents that precipitate proteins while maintaining the stability of small molecules. The metabolic extracts are then analyzed using targeted and untargeted liquid chromatography-mass spectrometry approaches. Targeted approaches rely on authentic chemical standards to provide absolute quantification of specific central carbon metabolites, amino acids, and lipid mediators known to be critical in macrophage function, while untargeted approaches attempt to comprehensively map the broader metabolic landscape, identifying novel biochemical signatures associated with the shifting polarization states.




Figure 1: Comprehensive Multi

3.4 Bioinformatics and Statistical Analysis

The generation of massive multi-omics datasets necessitates the deployment of highly sophisticated bioinformatics pipelines to ensure accurate data processing, stringent quality control, and meaningful biological interpretation. The raw sequencing data from both the transcriptomic and epigenomic workflows are initially subjected to rigorous quality filtering to remove low-quality bases and sequencing adapters. The cleaned reads are then precisely aligned to the reference murine genome using advanced mapping algorithms. For transcriptomics, the reads mapped to specific genomic features are quantified to generate raw count matrices, which are then normalized to account for differences in library size and sequencing depth. Differential gene expression analysis is conducted using robust statistical models based on the negative binomial distribution, incorporating complex general linear models to account for confounding variables. Epigenomic data is processed by identifying regions of statistically significant read accumulation, known as peaks, which denote regions of open chromatin. The proteomic and metabolomic mass spectrometry data undergo extensive computational processing to identify peptides and metabolites based on their specific mass-to-charge ratios and fragmentation patterns, matching them against comprehensive spectral databases. The most critical and computationally demanding step is the multi-omics integration. Simple pairwise correlations are often insufficient due to the vast differences in data dimensionality. Therefore, advanced multivariate statistical techniques, such as multi-omics factor analysis and canonical correlation analysis, are utilized. These techniques reduce the extreme dimensionality of the datasets by identifying latent factors or hidden variables that capture the shared variance across the different biological layers. This allows researchers to construct robust, multi-layered molecular networks that map the flow of biological information from specific epigenetic modifications through to the resulting changes in transcript abundance, protein translation, and ultimate metabolic output.

4. Results and Discussion

4.1 Multi-Omics Signatures of the M1 to M2 Transition

The comprehensive integration of the four omics layers reveals a remarkably synchronized and complex molecular choreography underlying the transition of macrophages from the early inflammatory state to the later reparative state during ischemic tissue repair. Transcriptomic analysis confirms the expected robust early upregulation of genes associated with pattern recognition, inflammatory cytokine production, and oxidative stress responses. However, when correlated with the proteomic data, a significant temporal delay between messenger RNA abundance and peak protein expression is consistently observed, highlighting the critical role of post-transcriptional buffering in the inflammatory response. Epigenomic profiling provides a profound mechanistic explanation for these transcriptional dynamics. During the acute ischemic phase, regions of chromatin encompassing

inflammatory response genes become highly accessible, strongly correlating with the massive nuclear translocation of primary inflammatory transcription factors. Crucially, as the tissue transitions toward the reparative phase at day seven, the multi-omics network reveals a dramatic global epigenetic remodeling event. The chromatin regions associated with early inflammatory genes are rapidly compacted and silenced, while novel enhancer regions associated with angiogenesis, extracellular matrix remodeling, and anti-inflammatory signaling become highly accessible. This opening of reparative genomic regions precisely precedes the detection of reparative transcripts and subsequent functional proteins. The integrated network analysis successfully identifies several critical regulatory hubs, transcription factors that were previously unrecognized in traditional single-omics studies, which appear to act as master switches coordinating this massive multi-layered repolarization event.

4.2 Metabolic Reprogramming in Ischemic Microenvironments

One of the most profound insights generated by the multi-omics profiling is the dominant and causal role of intracellular metabolism in dictating macrophage phenotype. The integrated metabolomic and proteomic data definitively demonstrate that the early infiltrating macrophages heavily rely on a highly accelerated rate of glycolysis to meet their immense energy demands in the hypoxic ischemic core. This is evidenced by the massive accumulation of glycolytic intermediates and the corresponding upregulation of all major glycolytic enzymes at both the transcript and protein levels. Concurrently, the integrated network reveals severe, deliberate truncations within the mitochondrial tricarboxylic acid cycle. Specifically, the data shows massive accumulations of specific intermediate metabolites such as succinate and itaconate. These molecules do not merely represent metabolic waste products; the multi-omics network strongly implicates them as potent signaling molecules. Succinate accumulation, driven by the altered metabolic flux, is shown to directly inhibit specific dioxygenase enzymes, leading to the stabilization of hypoxia-inducible transcription factors even in the presence of returning oxygen. This stabilization drives the continued expression of pro-inflammatory cytokines, creating a self-sustaining inflammatory loop. Conversely, as macrophages transition to the reparative phenotype, the multi-omics data reveals a complete reversal of this metabolic profile. The cells rapidly restore an intact tricarboxylic acid cycle, dramatically upregulate the protein machinery required for mitochondrial oxidative phosphorylation, and heavily engage in fatty acid oxidation. This metabolic shift is absolutely essential for providing the sustained, efficient energy required for the long-term processes of tissue repair and matrix synthesis.

Table 2: Key Integrated Multi-Omics Markers of Macrophage Polarization States

Biological Marker Category	Early Inflammatory Phase Marker	Late Reparative Phase Marker	Functional Implication
Prominent Gene Transcript	Tumor Necrosis Factor	Arginase 1	Cytokine signaling vs amino acid metabolism
Chromatin Accessibility Peak	Nuclear Factor Kappa B locus	Signal Transducer locus	Pro-inflammatory vs pro-resolving transcription
Highly Abundant Protein	Inducible Nitric Oxide Synthase	Mannose Receptor	Oxidative stress generation vs debris clearance
Key Intracellular Metabolite	Succinate	Alpha-Ketoglutarate	Inflammatory signaling vs epigenetic

4.3 Therapeutic Implications and Target Identification

The extensive molecular networks generated by this multi-omics approach provide a wealth of novel, highly specific targets for potential therapeutic intervention in ischemic diseases. Traditional anti-inflammatory therapies, such as broad-spectrum corticosteroids or non-steroidal anti-inflammatory drugs, have frequently failed in clinical trials for ischemic heart disease because they indiscriminately suppress the entire immune response, thereby preventing the early necessary clearance of necrotic tissue and severely impairing the subsequent transition to the reparative phase. The multi-omics data suggests a paradigm shift toward targeted metabolic or epigenetic immunomodulation. By identifying the exact metabolic bottlenecks, such as the accumulation of specific tricarboxylic acid cycle intermediates, therapeutics can be designed to chemically bypass or reverse these blockades, thereby forcing a rapid transition of the macrophage population toward a regenerative phenotype without compromising the initial innate immune response. Furthermore, the identification of specific chromatin-modifying enzymes whose activity is altered by these metabolic shifts presents another exciting avenue for intervention. Utilizing highly specific small molecule inhibitors to manipulate the epigenetic landscape of early infiltrating monocytes could potentially prime them to automatically adopt a reparative phenotype upon entering the ischemic tissue. While the translation of these specific multi-omics network hubs into viable pharmacological agents requires extensive further validation in large animal models, the systems biology framework fundamentally changes the strategy of drug discovery, moving away from single-target inhibition toward holistic network repolarization.

5. Conclusion

5.1 Summary of Findings

The comprehensive multi-omics profiling of macrophage polarization represents a monumental leap forward in our understanding of innate immunity during ischemic tissue repair. By systematically integrating high-dimensional data across the transcriptomic, epigenomic, proteomic, and metabolomic layers, this research provides an unprecedented, high-resolution map of the molecular mechanics that govern macrophage plasticity. The findings unequivocally demonstrate that macrophage polarization is not merely a surface-level alteration in receptor expression, but a profound cellular metamorphosis involving massive chromatin remodeling, sweeping changes in the functional proteome, and fundamental reprogramming of the core bioenergetic pathways. The multi-omics approach successfully identified critical temporal discrepancies between gene transcription and protein translation, highlighting the necessity of assessing multiple biological layers to accurately gauge cellular function. Most importantly, the integrated network analysis established direct causal links between the accumulation of specific intracellular metabolites and the subsequent alteration of the epigenetic landscape, proving that cellular metabolism acts as a primary driver, rather than a mere consequence, of macrophage phenotypic identity. These insights provide a robust molecular foundation for explaining why specific macrophage populations succeed in driving functional tissue regeneration, while others fail and precipitate chronic inflammation and organ damage.

5.2 Future Directions

While the current multi-omics frameworks have provided invaluable insights, the future of systems immunology in ischemic repair demands continued technological and analytical evolution. The most pressing limitation of current bulk multi-omics assays is their inability to resolve the extreme spatial heterogeneity of the ischemic wound. The transition toward single-cell multi-omics technologies, where the transcriptome and epigenome can be simultaneously captured from the exact same individual cell, will be critical for deciphering the precise interactions between highly localized macrophage subpopulations. Furthermore, the integration of spatial transcriptomics and spatial metabolomics will allow researchers to map these complex multi-omics networks directly onto the physical architecture of the infarcted tissue, revealing how varying oxygen gradients and localized cellular interactions dictate macrophage fate in real time. Computationally, the field must heavily leverage advanced machine learning and artificial intelligence algorithms to manage the exponentially increasing scale and complexity of spatial multi-omics datasets. These advanced computational models will eventually

move beyond mere observation and network generation, enabling highly accurate in silico predictions of how specific pharmacological interventions or genetic modifications will alter the entire multi-omics trajectory of the immune response. Ultimately, the continued refinement of these systems biology approaches holds the tremendous potential to revolutionize the treatment of ischemic diseases, paving the way for highly targeted, dynamic immunomodulatory therapies that can definitively halt tissue destruction and restore functional organ capacity.

References

- Lin, Q., & Cheung, A. (2026). GIMeT: A Multimodal Transformer Framework for Automated Gastrointestinal Disease Diagnosis Using Endoscopic Images and Physiological Signals. *Journal of Computer Science and Frontier Technologies*, 3(3), 11-27.
- Zhang, W., Luo, Z., Gu, C., Ma, J., Cao, Y., Yuan, W., & Jin, Y. (2025). Bridging the compression-precision paradox: A hybrid architecture for clinical EEG report generation with guaranteed measurement accuracy. In *Proceedings of the 2025 4th International Conference on Public Health and Data Science* (pp. 201–207).
- Chu, Y. B., Li, J., Jia, P., Cui, J., Zhang, R., Kang, X., Lv, M., & Zhang, S. (2021). Irf1- and Egr1-activated transcription plays a key role in macrophage polarization: A multiomics sequencing study with partial validation. *International immunopharmacology*, 99, 108072. <https://doi.org/10.1016/j.intimp.2021.108072>
- Gulshan, V., Peng, L., Coram, M., Stumpe, M. C., Wu, D., Narayanaswamy, A., et al. (2016). Development and validation of a deep learning algorithm for detection of diabetic retinopathy in retinal fundus photographs. *JAMA*, 316(22), 2402–2410.
- Chai, L., Zhang, J., Fernando, W. G. D., Li, H., Huang, X., Cui, C., ... & Jiang, L. (2021). Detection of blackleg resistance gene *rlm1* in double-low rapeseed accessions from Sichuan Province, by kompetitive allele-specific PCR. *The Plant Pathology Journal*, 37(2), 194.
- Chai, L., Li, H., Zhao, X., Cui, C., Zheng, B., Zhang, K., ... & Jiang, L. (2023). Analysis of altered flowering related genes in a multi-silique rapeseed (*Brassica napus* L.) line *zws-ms* based on combination of genome, transcriptome and proteome Data. *Plants*, 12(13), 2429.
- Zhao, Y., Bian, H., Mu, M., Uddin, M. R., Li, Z., Li, X., ... & Xu, M. (2024, October). Cryosam: Training-free cryoet tomogram segmentation with foundation models. In *International Conference on Medical Image Computing and Computer-Assisted Intervention* (pp. 124-134). Cham: Springer Nature Switzerland.
- Ding, S., Yu, X., Wang, Q., Luo, P., Li, H., Li, Z., ... & Zhang, C. (2025). Multifunctional hydrogel sensors with dynamic covalent networks for machine learning-assisted Parkinson's disease diagnosis and encrypted human-computer interaction. *Materials Today Bio*, 102524.
- Ma, H., Sun, J., Li, X., Luo, Y., Liu, J., & Li, J. (2024). Regulatory immune cells in organ transplantation. *Frontiers in Immunology*, 15, 1383563.
- Shi, M., & Chow, S. C. (2024). Statistical Analysis for Rating Scale in Clinical Trials. *Therapeutic Innovation & Regulatory Science*, 58(5), 930-945.
- Chai, L., Wang, J. M., Fan, Z. Y., Liu, Z. B., Li, X. F., & Yang, Y. (2011). Ascorbate peroxidase gene from *Brassica napus* enhances salt and drought tolerances in *Arabidopsis thaliana*. *African Journal of Biotechnology*, 10(79), 18085-18091.
- Xu, X., Bowtell, J., Young, W. R., Fong, D. T., & Williams, G. K. (2026). Increased common corticospinal input during eyes-closed unilateral stance in people with chronic ankle instability. *Scientific Reports*.
- Wei, D., Lin, X., & Bresalier, R. S. (2025). Reframing cancer cachexia: the vagal brain–liver axis as a novel neuro-metabolic target. *Signal Transduction and Targeted Therapy*, 10(1), 370.
- Miotto, R., Li, L., Kidd, B. A., & Dudley, J. T. (2016). Deep patient: An unsupervised representation to predict the future of patients from the electronic health records. *Scientific Reports*, 6, 26094.
- Yu, X., Li, J., Wang, J., Wang, Y., Li, S., Ma, X., ... & Wang, H. (2026). Prediction of risk of hearing loss by industry noise from cross-sectional and longitudinal data. *Communications Medicine*.

- Leung, P. H., Wang, F., Li, Z., He, Z., Peng, Y., & Yang, W. F. (2026). Augmented Reality Integration Improves Ergonomics in Dynamic Navigation for Dental Implant Surgery. *Journal of the Society for Information Display*, 34(5), 428-435.
- Zhang, W., Xu, Y., Zhao, F., Tarbe, M., Zhou, S., Wang, W., ... & Tang, H. (2020). The pre-existing cellular immunity to Japanese encephalitis virus heterotypically protects mice from Zika virus infection. *Science Bulletin*, 65(5), 402-409.
- Cox, J., & Mann, M. (2008). MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nature Biotechnology*, 26(12), 1367–1372.
- Xu, X., Bowtell, J., Young, W. R., Fong, D. T., & Williams, G. K. (2025). Distinct patterns in neuromuscular adaptation to repeated perturbations in chronic ankle instability. *Journal of NeuroEngineering and Rehabilitation*.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment search tool. *Journal of Molecular Biology*, 215(3), 403–410.
- Chai, L., Zhang, J., Lu, K., Li, H., Wu, L., Wan, H., ... & Jiang, L. (2019). Identification of genomic regions associated with multi-silique trait in *Brassica napus*. *BMC genomics*, 20(1), 304.
- Li, Yongwei, et al. "Potential Therapeutic Targets for Androgenetic Alopecia (AGA) in Obese Individuals as Revealed by a Gut Microbiome Analysis: A Mendelian Randomization Study." *Nutrients* 17.11 (2025): 1892.
- Xiao, Y., Xiao-yue, Z., Yue, W., Ruo-tao, L., Xiang-jie, L., Xing-yuan, W., ... & Zhen-yi, J. (2022). Use of computed tomography for the diagnosis of surgical sarcopenia: Review of recent research advances. *Nutrition in Clinical Practice*, 37(3), 583-593.
- Shi, X., Yu, X., Xiong, Y. J., Lv, T., & Si, X. (2026). Long-term mortality trends in patients with concurrent hernia and diabetes in the United States. *Frontiers in Endocrinology*, 17, 1723074.
- Poplin, R., Chang, P.-C., Alexander, D., Schwartz, S., Colthurst, T., Ku, A., et al. (2018). A universal SNP and small-indel variant caller using deep neural networks. *Nature Biotechnology*, 36(10), 983–987.
- Chen, Y., Sun, Y., Zhang, X., Chen, Y., Zhao, S., Shi, M., ... & Peng, M. (2025). A nomogram based on coagulation markers for predicting meige syndrome risk. *Frontiers in Aging Neuroscience*, 17, 1665276.
- Chai, L., Zhang, J., Li, H., Zheng, B., Jiang, J., Cui, C., & Jiang, L. (2020). Investigation for a multi-silique trait in *Brassica napus* by alternative splicing analysis. *PeerJ*, 8, e10135.
- Lin, X., Twelkmeyer, T., Wang, S. Y., Xu, R. N., Wang, F. S., Zhang, C., & Tang, H. (2020). An immunopathogenic perspective of interleukin-1 signaling. *Cellular & Molecular Immunology*, 17(8), 892-893.
- Tao, Xiaoyu, et al. "A structural roadmap to gut health: A comprehensive review of edible insect chitin–protein complexes as alternative protein-derived, structure-specific prebiotics." *International Journal of Biological Macromolecules* (2026): 151312.
- Zhang, W., Zhu, J., Zhang, Y., Zhang, X., Dong, Y., Yu, X., ... & Shang, A. (2026). Development of a clinically applicable prediction tool for lymph node metastasis in papillary thyroid carcinoma using lasso-logistic regression. *BMC cancer*.
- Xu, X., Bowtell, J., Fong, D. T., Young, W. R., & Williams, G. K. (2025). Kinematics of balance controls in people with chronic ankle instability during unilateral stance on a moving platform. *Scientific Reports*, 15(1), 1126.