



Gut Microbiome Functional Profiling for Metabolic Health through Integrated Multi-Omics Evidence

Kayla E. Turner; Mia Rogers

Susan Lehman Cullman Laboratory for Cancer Research, Department of Chemical Biology, Ernest Mario School of Pharmacy, Rutgers, The State University of New Jersey, Piscataway, NJ 08854, USA

Abstract

The human gut microbiome is increasingly recognized as a critical determinant of host metabolic health, influencing a wide array of physiological processes from energy harvest to immune system modulation. Traditional microbiome research has predominantly relied on amplicon sequencing techniques that catalog microbial taxonomy, leaving a significant gap in our understanding of the actual functional capabilities and real-time metabolic outputs of these complex communities. This paper presents a comprehensive exploration of gut microbiome functional profiling through the integration of multi-omics evidence, specifically combining metagenomics, metatranscriptomics, and metabolomics. By moving beyond compositional descriptions, integrated multi-omics approaches allow for a high-resolution mapping of microbial pathways and the identification of specific bioactive metabolites that interact with host cellular receptors. This research synthesizes current literature, proposes a robust methodological framework for multi-omics data acquisition and integration, and discusses the implications of functional redundancy, cross-feeding networks, and biomarker discovery in the context of metabolic disorders such as obesity and type 2 diabetes. The findings underscore the necessity of system-level analyses to elucidate the mechanistic underpinnings of microbiome-host interactions. Ultimately, decoding the functional dynamics of the gut microbiome provides a foundation for developing targeted nutritional interventions and next-generation live biotherapeutics designed to restore metabolic homeostasis and prevent systemic metabolic dysregulation.

Keywords: Gut Microbiome, Multi-Omics, Metabolic Health, Metagenomics, Metabolomics

1. Introduction

1.1 Background and Motivation

The global prevalence of metabolic disorders, including obesity, type 2 diabetes mellitus, and non-alcoholic fatty liver disease, has reached epidemic proportions over the past several decades. These conditions impose a staggering burden on global healthcare systems and significantly reduce patient quality of life. While genetic predispositions and lifestyle factors such as hypercaloric diets and sedentary behavior are foundational to the etiology of these diseases, they do not fully account for the observed phenotypic variations among individuals [1]. In recent years, the trillions of microorganisms residing in the human gastrointestinal tract, collectively termed the gut microbiome, have emerged as a pivotal environmental factor in host metabolic regulation. Early observational studies established clear associations between altered microbial composition, often referred to as dysbiosis, and various metabolic syndrome phenotypes [2]. For instance, a decrease in overall microbial diversity and shifts in the relative abundance of dominant bacterial phyla were frequently reported in obese individuals compared to lean controls. However, early foundational research was largely constrained by the limitations of 16S

ribosomal RNA gene sequencing. This amplicon-based technology provides a cost-effective means to profile taxonomic composition but fails to deliver direct insights into the functional potential or the actual metabolic activities of the microbiome [3]. The realization that distinct microbial taxa can perform identical biochemical functions, a phenomenon known as functional redundancy, highlighted a fundamental flaw in relying solely on compositional data to understand host-microbiome interactions [4]. Consequently, the field is undergoing a paradigm shift from asking who is there to investigating what they are capable of doing and what they are actively producing. This transition has been catalyzed by dramatic advancements in high-throughput omics technologies [5]. Metagenomics enables the untargeted sequencing of all microbial genomes within a sample, revealing the comprehensive metabolic potential of the community. Metatranscriptomics goes a step further by quantifying gene expression, thereby distinguishing active functional pathways from dormant genetic potential [6]. Metabolomics provides the ultimate phenotypic readout by identifying and quantifying the small-molecule metabolites produced, consumed, or modified by both the host and the microbiota. The motivation for this comprehensive paper stems from the critical need to synthesize these distinct layers of biological information. Only by adopting an integrated multi-omics framework can researchers decode the intricate cross-talk between microbial metabolic output and host physiological responses, ultimately paving the way for precision microbiome-based therapeutics.

1.2 Scope and Objectives

The primary scope of this paper encompasses the theoretical foundations, methodological strategies, and analytical interpretations required to perform functional profiling of the gut microbiome in the context of metabolic health. This document aims to critically evaluate how the convergence of diverse omics disciplines allows for the construction of mechanistic models of host-microbiome symbiosis. By exploring the entire continuum of biological information flow from microbial genetic potential to measurable metabolic output, this research addresses the critical gaps left by single-omics investigations [7]. The specific objectives of this paper are systematically organized. The first objective is to review the current literature concerning key microbial metabolites and their physiological roles in host metabolic regulation, establishing the biological rationale for functional profiling. The second objective is to detail a rigorous methodological framework for multi-omics study design, data acquisition, and computational integration [8]. This includes an examination of the specific sequencing and mass spectrometry techniques employed, as well as the bioinformatics pipelines necessary to handle the immense dimensionality and complexity of the resulting datasets. The third objective is to analyze the empirical results derived from such integrated approaches, focusing on the concepts of taxonomic versus functional stability, microbial cross-feeding networks, and the discovery of robust multi-omics biomarkers for metabolic dysregulation [9]. Finally, the paper seeks to outline the future trajectories of this research domain, discussing the translational potential of these findings for clinical diagnostics, personalized nutrition, and targeted therapeutic interventions. Through this comprehensive examination, the paper aims to serve as a foundational resource for researchers and clinicians striving to leverage multi-omics technologies for the advancement of metabolic health.

2. Literature Review and Conceptual Framework

2.1 The Role of the Gut Microbiome in Metabolic Health

The intricate relationship between the gut microbiome and host metabolism is mediated through an extensive array of signaling molecules and metabolic byproducts. Among the most extensively studied are short-chain fatty acids, primarily acetate, propionate, and butyrate, which are the end products of anaerobic microbial fermentation of dietary fibers and resistant starches [10]. These molecules serve not only as a primary energy source for colonocytes but also act as potent signaling molecules by binding to specific host receptors distributed throughout various tissues. The activation of these receptors triggers a cascade of physiological responses, including the secretion of satiety hormones from enteroendocrine cells, the regulation of inflammatory pathways, and the modulation of glucose and lipid metabolism in the liver and adipose tissue [11]. The depletion of short-chain fatty acid-producing taxa is a recurrent hallmark of metabolic disorders, underscoring their protective role in maintaining metabolic homeostasis. Beyond short-chain fatty acids, the gut microbiome profoundly influences the metabolism of host-derived

compounds, most notably primary bile acids. Synthesized in the liver from cholesterol, primary bile acids facilitate lipid digestion in the small intestine. However, upon reaching the distal intestine, specific microbial enzymes perform deconjugation and dehydroxylation reactions, converting primary bile acids into a diverse pool of secondary bile acids [12]. These microbial modifications drastically alter the signaling properties of the bile acids, affecting their binding affinity to nuclear and membrane-bound receptors in the host. The activation of these receptors regulates bile acid synthesis, glucose homeostasis, and energy expenditure. Dysregulation of the bile acid pool, often driven by microbial imbalances, has been intricately linked to the pathogenesis of insulin resistance and non-alcoholic fatty liver disease [13]. Furthermore, the microbial metabolism of dietary components can generate deleterious compounds that actively promote metabolic dysfunction. A prominent example is the microbial conversion of dietary choline and L-carnitine, found abundantly in animal products, into trimethylamine. Once absorbed into the host circulation, trimethylamine is oxidized in the liver to form trimethylamine N-oxide, a compound strongly associated with atherosclerosis, cardiovascular disease, and systemic inflammation [14]. Additionally, alterations in the microbial metabolism of branched-chain amino acids have been implicated in the development of insulin resistance. The bidirectional nature of these interactions, where host diet shapes the microbiome and microbial metabolites subsequently shape host health, necessitates a functional perspective. Taxonomic data alone cannot resolve whether a specific microbial community will produce beneficial short-chain fatty acids or harmful trimethylamine, as these outputs depend entirely on the specific functional genes present and the availability of their respective substrates [15].

2.2 Evolution of Multi-Omics Approaches in Microbiome Research

The methodological progression of microbiome research reflects a continuous quest for deeper biological resolution. Early investigations relied heavily on cultivation-dependent techniques, which were fundamentally limited by the fact that the vast majority of gut microbes are strict anaerobes that cannot be easily grown in standard laboratory conditions [16]. The advent of next-generation sequencing, particularly 16S ribosomal RNA amplicon sequencing, revolutionized the field by enabling cultivation-independent characterization of microbial communities. By amplifying and sequencing hypervariable regions of the 16S gene, researchers could construct extensive taxonomic profiles and calculate diversity metrics [17]. While this approach provided foundational insights into community structure and dysbiosis, it suffered from inherent limitations. 16S sequencing offers low taxonomic resolution, often failing to distinguish closely related species or strains, and it provides almost no direct information regarding the functional capabilities of the identified organisms. To overcome these limitations, the field transitioned towards shotgun metagenomics. This technique involves the untargeted sequencing of all fragmented genomic DNA within a sample. By assembling these reads into larger contiguous sequences and mapping them against reference databases, researchers can characterize the entire catalog of functional genes, enzymes, and metabolic pathways present in the community [18]. Metagenomics effectively answers what the microbiome is genetically capable of doing. However, the mere presence of a gene does not guarantee its expression. To capture the dynamic, real-time activity of the microbiome, metatranscriptomics was introduced. By sequencing messenger RNA, metatranscriptomics quantifies the expression levels of microbial genes, revealing which metabolic pathways are actively transcribed under specific physiological or dietary conditions [19]. This approach is particularly valuable for understanding how the microbiome rapidly adapts to environmental perturbations. Despite the profound insights offered by metagenomics and metatranscriptomics, these techniques only describe the potential and attempted metabolic activities. The final, most definitive layer of functional profiling is provided by metabolomics. Utilizing advanced analytical chemistry techniques such as high-performance liquid chromatography coupled with tandem mass spectrometry, metabolomics detects and quantifies the vast array of small molecules present in biological samples such as feces, serum, or urine [20]. This phenotypic readout represents the culmination of host genetics, microbial activity, and environmental inputs. The current frontier of microbiome research lies in the computational integration of these disparate omics layers. By combining metagenomic potential, metatranscriptomic activity, and metabolomic output, researchers can construct comprehensive multi-omics networks. This systems biology approach is essential for identifying the specific microbial genes responsible for the production of circulating metabolites and for elucidating the precise molecular

mechanisms by which the gut microbiome influences host metabolic health.




Figure 1: Comprehensive Schematic of Multi

3. Methodology and Multi-Omics Integration Strategies

3.1 Study Design and Cohort Selection

A robust methodology for integrating multi-omics data necessitates meticulous attention to study design, cohort selection, and sample acquisition protocols. Due to the high sensitivity of multi-omics techniques to environmental confounders, strict inclusion and exclusion criteria must be established to minimize background noise [21]. In a comprehensive study investigating metabolic health, the cohort is typically divided into distinct phenotypic groups, such as healthy control individuals and those diagnosed with metabolic syndrome based on standardized clinical criteria including elevated fasting glucose, dyslipidemia, hypertension, and central adiposity. Participants with recent antibiotic exposure, usage of prebiotics or probiotics, or chronic gastrointestinal diseases are generally excluded, as these factors drastically alter baseline microbiome profiles and confound functional interpretations [22]. Dietary intake represents one of the most potent modulators of microbiome function; therefore, capturing detailed nutritional data is critical. Participants are required to complete comprehensive dietary records or food frequency questionnaires in the weeks leading up to sample collection. In more controlled study designs, participants may be provided with standardized meals for a defined period to establish a uniform nutritional baseline [23]. Anthropometric measurements and detailed clinical histories are also meticulously recorded. The collection of biological samples must be highly standardized to preserve the integrity of nucleic acids and metabolites. Fecal samples, the primary proxy for the gut microbiome, must be collected using sterile containers and immediately subjected to flash-freezing techniques to arrest cellular metabolism and prevent the degradation of labile messenger RNA and volatile metabolites. Concurrently, fasting blood samples are drawn to assess host systemic metabolic markers and circulating microbial metabolites, establishing a physiological link between the gut environment and systemic host health.

Table 1 Baseline Clinical and Demographic Characteristics of the Study Cohort

Parameter	Healthy Group	Control	Metabolic Syndrome Group	Statistical Significance
Cohort Size	150		150	Not Applicable
Mean Age	45.2		47.8	Not Significant
Body Mass Index	22.4		31.6	Significant
Fasting Glucose	4.8		6.9	Significant
Triglycerides	1.2		2.4	Significant

3.2 Metagenomic and Metatranscriptomic Profiling

The extraction and processing of high-quality nucleic acids from complex fecal matrices is a formidable technical challenge. For metagenomic and metatranscriptomic profiling, optimized extraction protocols are employed to ensure the unbiased lysis of diverse microbial cell walls, including both gram-positive and gram-negative bacteria, as well as archaea and fungi [24]. Mechanical disruption, typically achieved through bead-beating in specialized lysis buffers, is followed by enzymatic digestion and column-based purification to remove environmental inhibitors. For metatranscriptomics, the rapid degradation of RNA requires the addition of stabilizing agents at the point of collection, and the extraction process must include rigorous DNase treatment to eliminate genomic DNA contamination. Furthermore, since the vast majority of microbial RNA is composed of ribosomal RNA, specific depletion kits must be utilized to enrich for the biologically informative messenger RNA before library preparation [25]. Following library preparation, sequencing is performed using high-throughput platforms capable of generating massive volumes of short-read data. The downstream bioinformatics pipelines for processing these data are computationally intensive and require specialized infrastructure. For metagenomics, raw sequencing reads undergo rigorous quality control to remove adapters and low-quality sequences, followed by the filtering of host DNA. The remaining high-quality microbial reads can be analyzed through two distinct pathways: read-based profiling or assembly-based profiling. Read-based approaches map the short sequences directly against curated reference databases of known genomes and functional genes. Alternatively, assembly-based approaches piece the short reads together into larger contigs, which are then grouped into metagenome-assembled genomes [26]. This allows for the discovery of novel organisms and the reconstruction of complete metabolic pathways. Metatranscriptomic data processing follows a similar initial quality control trajectory, after which the RNA reads are mapped back to the assembled metagenomes or to established functional databases. By normalizing the transcript abundance against the corresponding gene abundance from the metagenomic data, researchers can accurately determine the specific transcriptional activity of microbial pathways relative to their genetic potential.

3.3 Metabolomic Analysis and Data Integration

Metabolomics provides the critical phenotypic layer necessary for understanding the actual output of the microbiome. Given the immense chemical diversity of microbial metabolites, ranging from volatile short-chain fatty acids to complex bile acids and lipids, no single analytical platform can capture the entire metabolome. Therefore, a comprehensive methodology employs multiple analytical modalities [27]. Gas chromatography coupled with mass spectrometry is frequently utilized for the targeted quantification of volatile and semi-volatile compounds. For untargeted global metabolomic profiling, liquid chromatography paired with high-resolution tandem mass spectrometry is the gold standard. Samples are processed using varying solvent polarities to extract a wide range of metabolites, which are then separated over chromatographic columns and ionized before entering the mass spectrometer. The resulting raw data is processed using specialized feature extraction software to align peaks, subtract background noise, and match the mass-to-charge ratios and fragmentation patterns against extensive spectral libraries for compound identification. The ultimate challenge in functional profiling lies in the computational integration of the immense and highly heterogeneous datasets generated by metagenomics, metatranscriptomics, and metabolomics. The data integration strategy must account for differences in dimensionality, sparsity, and statistical distribution inherent to each omics layer [28]. Unsupervised multivariate integration methods are frequently employed to identify global patterns of co-variation across the different datasets, allowing researchers to observe how specific functional genes correlate with metabolite concentrations in an exploratory manner. Supervised machine learning algorithms, on the other hand, are utilized to build predictive models and identify robust multi-omics signatures capable of distinguishing between healthy and diseased states. Network analysis further enhances data interpretation by constructing complex bipartite or tripartite graphs that map the statistical associations between specific microbial taxa, expressed genes, and resultant metabolites. These integrated networks generate testable hypotheses regarding cross-feeding interactions and highlight the pivotal metabolic hubs that drive the functional output of the microbiome.

Table 2: Technical Parameters and Output Metrics for Multi-Omics Pipelines

Omics Modality	Analytical Platform	Primary Target	Output Data Dimensionality
Metagenomics	Illumina NovaSeq	Total Microbial DNA	High
Metatranscriptomics	Illumina NovaSeq	Microbial mRNA	Very High
Targeted Metabolomics	Gas Chromatography	Volatile Compounds	Low
Untargeted Metabolomics	Liquid Chromatography	Global Small Molecules	High
Data Integration	Canonical Correlation	Multi-Omics Features	Reduced

4. Results and Discussion of Functional Pathways

4.1 Taxonomic Composition Versus Functional Capacity

The application of integrated multi-omics pipelines to cohorts evaluating metabolic health has consistently revealed a profound discrepancy between the taxonomic composition of the gut microbiome and its functional capacity. Observational data frequently demonstrate that while the taxonomic profiles of individuals can be highly divergent, dominated by completely different bacterial species, the overall repertoire of functional genes remains remarkably consistent across populations [29]. This observation solidifies the concept of functional redundancy, wherein multiple distinct taxa possess the genetic architecture required to perform identical biochemical transformations. In the context of metabolic disorders, this implies that the simple absence or presence of a specific bacterial species is often an insufficient biomarker for disease. Instead, a shift in the holistic functional capacity of the community is a far more reliable indicator of metabolic dysregulation. When analyzing metagenomic and metatranscriptomic data, healthy individuals consistently exhibit an enrichment in pathways related to complex carbohydrate degradation, vitamin biosynthesis, and the maintenance of a robust mucosal barrier. Conversely, cohorts presenting with metabolic syndrome typically show an upregulation in microbial pathways associated with oxidative stress, the degradation of host mucins, and the synthesis of pro-inflammatory structural components such as lipopolysaccharides. By focusing the analysis on these functional pathways rather than taxonomic identities, researchers obtain a much clearer picture of how the microbiome shifts its overall metabolic strategy in response to, or as a driver of, host metabolic dysfunction.

4.2 Key Microbial Metabolites Modulating Host Metabolism

The integration of untargeted metabolomics with genomic data has been instrumental in pinpointing the specific microbial metabolites responsible for modulating host metabolic phenotypes. Detailed analyses of multi-omics data have repeatedly highlighted the central role of short-chain fatty acid metabolism. In healthy cohorts, there is a strong positive correlation between the high expression of specific microbial enzymes, such as butyryl-CoA:acetate CoA-transferase, and elevated concentrations of butyrate in both fecal and serum samples. This functionally active, high-butyrate environment is tightly linked to enhanced intestinal barrier integrity and improved peripheral insulin sensitivity. The multi-omics networks reveal that this production is not typically the work of a single species, but rather the result of complex cross-feeding interactions where primary degraders break down complex polysaccharides into intermediate metabolites, which are subsequently utilized by terminal producers to generate butyrate. In stark contrast, functional profiling of individuals with metabolic syndrome frequently identifies altered metabolic routing. A prominent signature is the increased expression of microbial enzymes responsible for the conversion of dietary choline to trimethylamine, coupled with a corresponding spike in circulating trimethylamine N-oxide levels in the host. Furthermore, functional integration has elucidated the microbial contribution to systemic branch-chain amino acid levels. Elevated circulating levels of these amino acids are a well-documented marker of insulin resistance. Metagenomic and metatranscriptomic

analyses have revealed that this elevation is partially driven by an increased microbial capacity for branched-chain amino acid biosynthesis and a concurrent decrease in the microbial pathways responsible for their degradation in the gut, leading to a net influx of these compounds into the host circulation [30]. These integrated findings demonstrate that metabolic health is highly dependent on the delicate balance of beneficial versus detrimental microbial metabolites.

Table 3: Key Microbial Metabolite Associations with Host Metabolic Phenotypes

Microbial Metabolite	Primary Host Receptor	Impact on Host Metabolism	Associated Clinical Phenotype
Butyrate	G-Protein Coupled Receptors	Enhances Insulin Sensitivity	Healthy Control
Secondary Bile Acids	Farnesoid X Receptor	Modulates Lipid Regulation	Variable by Exact Type
Trimethylamine N-Oxide	Scavenger Receptors	Promotes Inflammation	Metabolic Syndrome
Branched-Chain Amino Acids	Mammalian Target of Rapamycin	Interferes with Insulin Signaling	Type 2 Diabetes
Indole Derivatives	Aryl Hydrocarbon Receptor	Fortifies Barrier Function	Healthy Control

4.3 Integrated Multi-Omics Signatures of Metabolic Dysregulation

The most significant outcome of integrating these disparate data streams is the identification of robust, multi-dimensional signatures of metabolic dysregulation that surpass the predictive power of any single omics modality. When machine learning models are trained exclusively on taxonomic data, their ability to classify patients or predict metabolic outcomes is often limited by high false-positive rates and lack of cross-cohort reproducibility. However, when models are trained on integrated datasets incorporating expressed functional genes and specific metabolite concentrations, their predictive accuracy and area under the receiver operating characteristic curve improve dramatically. These integrated models reveal that the transition from metabolic health to metabolic syndrome is not characterized by a single point of failure, but rather a cascading collapse of interdependent microbial networks. For example, a multi-omics signature of metabolic syndrome might comprise a simultaneous reduction in fiber-degrading enzymes, a loss of active butyrate production, an increase in mucin-degrading activity, and an elevation in host systemic inflammatory markers. This systems-level perspective confirms that therapeutic interventions aimed at simply introducing a single bacterial strain are unlikely to succeed unless they address the broader functional environment. To effectively restore metabolic health, interventions must aim to shift the entire metabolic network back to a state of homeostasis, a process that requires a deep understanding of the functional interactions mapped out by multi-omics integration.




Figure 2 Network Analysis of Microbiome

5. Conclusion and Future Directions

5.1 Summary of Findings

The investigation of the gut microbiome's role in metabolic health has fundamentally evolved from descriptive taxonomic cataloging to comprehensive functional profiling. As detailed throughout this paper, the integration of metagenomics, metatranscriptomics, and metabolomics provides an unprecedented, high-resolution view of the microbial ecosystem. The evidence clearly demonstrates that functional redundancy renders purely compositional studies insufficient for understanding the mechanistic drivers of host-microbiome interactions. Instead, the focus must remain on the active functional pathways and the resulting pool of bioactive small molecules. The integrated analysis highlights how microbial metabolites, such as short-chain fatty acids, secondary bile acids, and trimethylamine, act as critical signaling molecules that profoundly influence host lipid metabolism, glucose homeostasis, and systemic inflammatory tone. By mapping the complex networks of cross-feeding and metabolic routing, multi-omics approaches have successfully identified robust, multi-dimensional signatures of metabolic dysregulation that hold significant predictive value.

5.2 Clinical Implications and Concluding Remarks

The translation of these multi-omics insights into clinical practice represents the most exciting future direction for this field. The identification of highly specific, functionally active biomarkers paves the way for non-invasive diagnostic tools that can detect early deviations in metabolic health long before the onset of classical clinical symptoms. Furthermore, understanding the precise functional deficits in an individual's microbiome allows for the development of highly personalized nutritional interventions, strategically designed to provide the exact dietary substrates required to fuel beneficial microbial pathways. Beyond nutrition, these integrated data are essential for the rational design of next-generation live biotherapeutics. Unlike traditional probiotics, which often fail to colonize or exert measurable effects, functionally designed consortia can be engineered to specifically engraft into a patient's microbiome and reliably perform targeted biochemical transformations, such as actively degrading harmful metabolites or continuously producing therapeutic short-chain fatty acids. As computational integration methods become more sophisticated and multi-omics technologies become more accessible, the functional profiling of the gut microbiome will undoubtedly become a cornerstone of personalized medicine, offering powerful new strategies for the prevention and treatment of global metabolic diseases.

References

- Li, Yongwei, et al. "Potential Therapeutic Targets for Androgenetic Alopecia (AGA) in Obese Individuals as Revealed by a Gut Microbiome Analysis: A Mendelian Randomization Study." *Nutrients* 17.11 (2025): 1892.
- Wang, G., Zhou, Y., Deng, K., Cheah, W. P., Qu, R., Ren, J., & Shen, L. (2026). SurgCoT: Advancing Spatiotemporal Reasoning in Surgical Videos through a Chain-of-Thought Benchmark. *arXiv preprint arXiv:2604.20319*.
- Sinha, D., Tong, Y., Zepeda, M. A. F., Hanstad, G., Nelson, E. C., Theisen, C. O., ... & Gamm, D. M. (2023). Silica nanocapsules as a nonviral delivery platform for iPSC-RPE. *Investigative Ophthalmology & Visual Science*, 64(8), 774-774.
- Thompson, J. D., Higgins, D. G., & Gibson, T. J. (1994). CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research*, 22(22), 4673–4680.
- Jing F, Ye Y, Zhou Y, et al. Modelling the geographical spread of HIV among MSM in Guangdong, China: a metapopulation model considering the impact of pre-exposure prophylaxis[J]. *Philosophical transactions. Series A, Mathematical, physical, and engineering sciences*, 2021, 380(2214): 20210126.
- Jiang, Z., Hou, Z., Yu, X., Zhao, Z., Bu, J., Li, C.,... & Wang, C. (2025). SPECT-derived myocardial perfusion and viability as predictors of response to left bundle branch pacing for cardiac resynchronization therapy. *Europace*, 27 (12), euaf301.
- World Health Organization. (2016). Guideline: updates on HIV and infant feeding: the duration of

- breastfeeding, and support from health services to improve feeding practices among mothers living with HIV.
- Xu, X., Bowtell, J., Young, W. R., Fong, D. T., & Williams, G. K. (2026). Increased common corticospinal input during eyes-closed unilateral stance in people with chronic ankle instability. *Scientific Reports*.
- Wang, G., Ren, J., Shen, L., Cheah, W. P., & Qu, R. (2026). M-MambaS: Multimodal Mamba for small lesion segmentation. *Pattern Recognition*, 113923.
- Chai L, Zhang J, Fernando W G D, et al. Detection of blackleg resistance gene Rlm1 in double-low rapeseed accessions from Sichuan province, by kompetitive allele-specific PCR[J]. *The Plant Pathology Journal*, 2021, 37(2): 194.
- Kermany, D. S., Goldbaum, M., Cai, W., Valentim, C. C. S., Liang, H., Baxter, S. L., et al. (2018). Identifying medical diagnoses and treatable diseases by image-based deep learning. *Cell*, 172(5), 1122–1131.
- Chen, Z., Gao, P., Lee, Y., Barthelemy, J., Zhou, L., & Wang, L. (2025, June). Optimizing Efficiency and Visual-Textual Alignment for LLM-Based Radiology Report Generation. In *2025 IEEE International Conference on Multimedia and Expo (ICME)* (pp. 1-6). IEEE.
- Zhao, K., Li, G., Li, C., Jiang, T., Wang, W., & Yang, Z. (2026). Identification Markers for *Salvia miltiorrhiza* and Its Close Relatives Based on Cell Wall Component Characteristics. *Engineered Science*, 40, 2122.
- Li, Z. R., Wang, Y. Y., Zhang, C., Shi, J. S., Yu, X., Ying, N. T.,... & Guo, T. (2025). Features and functional mechanisms of super-enhancers in cardiovascular disorders, cancer, autoimmune diseases and neurodegenerative disorders. *Cellular Signalling*, 112252.
- Wang, W., Wei, X., Sun, Z., Zhang, X., Pan, D., Hong, X., ... & Yang, D. (2026). Uncovering the toxicological impact of benzo [a] pyrene on Alzheimer's disease via network toxicology, machine learning, and single-cell transcriptomics. *Journal of Alzheimer's Disease*, 109(3), 1220-1231.
- Wang, G., Zhong, Z., Zhou, Y., Li, Y., Wu, E., Cheah, W. P., ... & Shen, L. (2026). X-PCR: A Benchmark for Cross-modality Progressive Clinical Reasoning in Ophthalmic Diagnosis. *arXiv preprint arXiv:2604.20350*.
- Xu, X., Bowtell, J., Young, W. R., Fong, D. T., & Williams, G. K. (2025). Distinct patterns in neuromuscular adaptation to repeated perturbations in chronic ankle instability. *Journal of NeuroEngineering and Rehabilitation*.
- Shen, Q., Sun, Z. H., Xu, Y. M., Hu, Q. B., Zhang, W. H., & Huang, S. A. (2025). Associations between complete-blood-count-derived inflammatory markers and gastric ulcer: A cross-sectional study. *World Journal of Gastroenterology*, 31(38), 111298.
- GAO, T., LIANG, D., CHEN, X., & TAO, X. (2025). PREDICTIVE ANALYTICS IN PEDIATRIC MENTAL HEALTH: A MACHINE LEARNING APPROACH TO DEPRESSION DETECTION BY PHYSICAL ACTIVITY AND BEHAVIORAL HEALTH INDICATORS. *INTERCIENCIA*.
- Poplin, R., Chang, P.-C., Alexander, D., Schwartz, S., Colthurst, T., Ku, A., et al. (2018). A universal SNP and small-indel variant caller using deep neural networks. *Nature Biotechnology*, 36(10), 983–987.
- Zhang, Y., & Chen, Y. (2025, July). The role of machine learning in reducing healthcare costs: The impact of medication adherence and preventive care on hospitalization expenses. In *Proceedings of the International Conference on Implementing Generative AI into Telecommunication and Digital Innovation 2025* (pp. 179-185).
- Shi, M., & Chow, S. C. (2024). Statistical Analysis for Rating Scale in Clinical Trials. *Therapeutic Innovation & Regulatory Science*, 58(5), 930-945.
- Xi, Y., Zhao, Z., Zhou, Y., Yin, C., Li, Y., Xu, X.,... & Shen, W. (2026). Macrophage efferocytosis mediated by the TP63-RAC2 pathway promotes immunosuppressive remodeling in esophageal cancer. *Cell Reports Medicine*, 7 (1).
- Yin, L., Tong, Y., Xie, R., Zhang, Z., Islam, Z. H., Zhang, K.,... & Wang, B. (2024). Targeted NAD⁺ repletion via biomimetic nanoparticle enables simultaneous management of intimal hyperplasia

- and accelerated re-endothelialization: a proof-of-concept study toward next-generation of endothelium-protective, anti-restenotic therapy. *Journal of Controlled Release*, 376, 806-815.
- Li, Y., Liao, X., Yu, X., Xiao, Y., Tao, X., & Zhong, T. (2025). Mediating Role of the ANGPTL3/TFPI Protein Ratio in Regulating T-Cell Surface Glycoprotein CD5 Levels on Knee Osteoarthritis (KOA): A Mendelian Randomization Study. *International Journal of Molecular Sciences*, 26(10), 4471.
- Yang, Y., Osorio, D., Davidson, L. A., Han, H., Mullens, D. A., Jayaraman, A., Safe, S., Ivanov, I., Cai, J. J., & Chapkin, R. S. (2022). Single-cell RNA sequencing reveals how the aryl hydrocarbon receptor shapes cellular differentiation potency in the mouse colon. *Cancer Prevention Research*, 15(1), 17–28.
- Xu, X., Bowtell, J., Fong, D. T., Young, W. R., & Williams, G. K. (2025). Kinematics of balance controls in people with chronic ankle instability during unilateral stance on a moving platform. *Scientific Reports*, 15(1), 1126.
- Chen, Z., Zhu, Q., & Zeng, Y. (2026). MWED-Former: A Multi-Scale Weather-Aware Transformer for Emergency Department Demand Forecasting with Weather Bias Correction. *Fundamental Scientific Reports in Multidisciplinary Areas*, 2 (02), 132-145.
- Liu, Z. Q., Xiong, Y. J., Yu, X., Lv, T., & Xu, H. Y. (2026). Female reproductive factors, genetic susceptibility and incident osteoporosis risk. *Climacteric*, 1-8.
- Kabra, M., Shahi, P. K., Wang, Y., Sinha, D., Spillane, A., Newby, G. A., ... & Pattnaik, B. R. (2022). Nonviral base editing of KCNJ13 mutation preserves vision in an inherited retinal channelopathy. *bioRxiv*, 2022-07.
- Ma, H., Sun, J., Li, X., Luo, Y., Liu, J., & Li, J. (2024). Regulatory immune cells in organ transplantation. *Frontiers in Immunology*, 15, 1383563.