



Ion-Specific Free-Energy Landscapes of Graphene-Lipid Bicelle Nucleation

Daniel Álvarez Ortega

Department of Mechanical Engineering, Faculty of Engineering, Universidad de Buenos Aires, Buenos Aires, Buenos Aires F.D., Argentina

Abstract

The interaction between carbon-based nanomaterials and biological membranes represents a critical frontier in nanomedicine, nanotoxicology, and bioengineering. Among these materials, pristine and functionalized graphene exhibit unique physicochemical properties that drive complex interfacial behaviors when exposed to lipid assemblies. This study presents a comprehensive investigation into the ion-specific free-energy landscapes governing the nucleation and structural evolution of graphene-lipid bicelles. Utilizing advanced computational frameworks, this research systematically explores how different monovalent and divalent cations dictate the thermodynamic pathways of lipid self-assembly on graphene surfaces. The findings reveal that ion specificity significantly alters the hydration dynamics, interfacial electrostatic screening, and entropic contributions during the nucleation process. Divalent cations such as calcium and magnesium are shown to induce profound structural stabilization of the nascent bicelle through ion-bridging mechanisms, whereas monovalent ions primarily modulate the solvent structure and hydrophobic driving forces in accordance with their respective positions in the Hofmeister series. The extracted free-energy profiles provide quantitative insights into the activation barriers and metastable states characteristic of these self-assembling systems. These insights offer fundamental guidelines for designing graphene-based nanoscale delivery vehicles and evaluating their potential cyto-compatibility in complex physiological fluid environments.

Keywords: Graphene, Lipid Bicelle, Ion Specificity, Free-Energy Landscape

1. Introduction

1.1 Background on Graphene and Lipid Interactions

The rapid advancement of nanotechnology has positioned graphene and its structural derivatives at the forefront of biomedical applications, ranging from targeted drug delivery vectors and biosensors to tissue engineering scaffolds. The unique two-dimensional atomic structure of graphene, characterized by exceptional mechanical strength, electrical conductivity, and high surface area, makes it an ideal candidate for interfacing with biological systems. However, the translation of graphene-based technologies into clinical settings is heavily contingent upon a rigorous understanding of their interactions with cellular boundaries, primarily the lipid bilayer membrane. When graphene sheets are introduced into a physiological environment, they immediately encounter a complex milieu of lipids, proteins, and electrolytes. The spontaneous adsorption of lipid molecules onto the hydrophobic basal plane of graphene initiates a self-assembly process that can lead to the formation of hybrid nanostructures, such as lipid-coated graphene or graphene-lipid bicelles. These bicelles are planar, disc-shaped lipid assemblies that serve as excellent biomimetic models for cell membranes, offering structural stability and retaining the lateral mobility of natural lipid bilayers. Understanding the mechanistic details of how these bicelles nucleate and grow on a graphene template is paramount for

predicting the biological fate, potential toxicity, and therapeutic efficacy of these nanomaterials. Recent investigations have demonstrated that the driving forces behind this self-assembly are predominantly hydrophobic interactions between the lipid acyl chains and the non-polar graphene surface, coupled with the complex interplay of solvent dynamics [1]. Nevertheless, the precise thermodynamic pathways and the transient intermediate states involved in the initial nucleation phase remain largely obscured by the rapid temporal scales and minute spatial dimensions characteristic of these interfacial phenomena [2].

1.2 The Role of Ion Specificity in Self-Assembly

Biological fluids are essentially complex aqueous electrolyte solutions containing a variety of dissolved ions, including sodium, potassium, calcium, magnesium, and chloride. These ions do not merely act as passive bystanders that screen electrostatic interactions; rather, they actively participate in modulating the structure and thermodynamics of biomolecular self-assembly. The phenomenon whereby different ions exert distinct effects on the physical properties of macromolecules and interfaces is broadly categorized under specific ion effects, often following the historical Hofmeister series. In the context of graphene-lipid bicelle nucleation, the local ionic environment plays a transformative role. The hydration free energy of the ions, their propensity to accumulate at or deplete from the liquid-solid interface, and their specific interactions with the zwitterionic or charged headgroups of the lipid molecules collectively dictate the overarching free-energy landscape of the system. For instance, strongly hydrated kosmotropic ions and weakly hydrated chaotropic ions perturb the hydrogen-bonding network of interfacial water molecules to varying degrees, thereby altering the hydrophobic effect that drives lipid adsorption onto the graphene substrate [3]. Furthermore, multivalent ions such as calcium have been shown to coordinate strongly with the phosphate moieties of phospholipid headgroups, leading to structural dehydration and the formation of highly stable ion-bridged complexes [4]. These ion-specific mechanisms suggest that the energy barriers for bicelle nucleation, as well as the morphological characteristics of the resulting graphene-lipid hybrids, are exquisitely sensitive to the background electrolyte composition. Despite the recognized importance of these effects, there is a notable gap in quantitative thermodynamic studies that systematically map the free-energy landscapes of these assemblies across a spectrum of biologically relevant ionic species [5].

1.3 Research Objectives and Scope

This paper aims to bridge the aforementioned knowledge gap by providing a detailed, quantitative analysis of the ion-specific free-energy landscapes associated with graphene-lipid bicelle nucleation. The primary objective is to elucidate how variations in cation identity and valency modulate the thermodynamic driving forces, activation barriers, and stable intermediate states during the self-assembly of dimyristoylphosphatidylcholine lipid molecules onto a pristine graphene nanosheet. By isolating the effects of sodium, potassium, magnesium, and calcium ions, this research systematically dissects the relative contributions of electrostatic screening, ion-lipid coordination, and solvent structuring to the overall nucleation process. The scope of this study encompasses extensive computational modeling and thermodynamic analysis, utilizing advanced enhanced sampling techniques to reconstruct the free-energy surfaces along chosen reaction coordinates. The subsequent sections will detail the theoretical underpinnings of specific ion effects at hydrophobic interfaces, describe the rigorous methodological framework employed to extract the free-energy profiles, and present a comprehensive discussion of the structural and energetic consequences of ion specificity. Ultimately, this work seeks to establish a predictive mechanistic understanding of nanomaterial-lipid interactions, providing essential theoretical guidelines for the rational design of safe and effective graphene-based nanomedicines tailored for specific physiological environments.

2. Literature Review and Theoretical Background

2.1 Nanomaterial-Biological Interface Dynamics

The interaction between engineered nanomaterials and biological constituents forms the crux of the rapidly evolving field of bio-nanotechnology. At the heart of this discipline lies the concept of the nano-bio interface, a dynamic and highly complex boundary where synthetic surfaces meet the molecular machinery of life. Historically, the evaluation of nanomaterial biocompatibility was largely

phenomenological, relying on macroscopic assays to determine cellular viability or tissue-level inflammation. However, a paradigm shift has occurred over the last decade, emphasizing the necessity of understanding these interactions at the fundamental atomic and molecular levels [6]. Graphene, owing to its prototypical two-dimensional geometry and profound hydrophobicity, has served as a central model system in this theoretical advancement. When a pristine graphene sheet contacts a biological membrane, it induces significant structural perturbations. Studies have shown that the strong van der Waals dispersion forces between the carbon lattice and the hydrocarbon tails of lipid molecules can overcome the cohesive forces holding the lipid bilayer together, leading to phenomena such as lipid extraction, membrane thinning, and pore formation [7]. Conversely, under controlled conditions, lipid molecules can cooperatively assemble onto the graphene surface to form stable, biologically inert coatings that mitigate toxicity. The kinetics and thermodynamics of this phase transition from dispersed monomeric lipids or spherical liposomes to flat, substrate-supported bicelles are governed by a delicate balance of enthalpic interactions and entropic penalties [8]. Enthalpically, the system seeks to maximize the contact between the lipid acyl chains and the graphene surface while minimizing the exposure of hydrophobic regions to the aqueous solvent. Entropically, the self-assembly imposes severe translational and conformational restrictions on the lipid molecules, which must be offset by the release of structured interfacial water into the bulk solution. This delicate thermodynamic balance renders the nucleation process highly susceptible to subtle environmental perturbations, particularly variations in the composition of the surrounding fluid [9].

2.2 Lipid Bicelles and Membrane Models

To systematically investigate the complexities of the nano-bio interface, researchers rely on robust and well-defined model membrane systems. While spherical liposomes and solid-supported lipid bilayers have been extensively utilized, lipid bicelles have emerged as particularly advantageous models for studying interactions with two-dimensional nanomaterials [10]. Bicelles are disk-shaped, self-assembled structures composed of a planar central region formed by long-chain phospholipids, bordered by a highly curved rim composed of short-chain lipids or specific detergents. This unique geometry effectively segregates the planar bilayer properties from the highly curved edge defects, providing a pristine, flat biological surface that closely mimics the lateral organization and fluidity of intact cellular membranes without the topological constraints of spherical vesicles. When interacting with graphene, the planar region of the bicelle exhibits a high degree of structural conformity with the flat carbon lattice. The nucleation of a bicelle on graphene is essentially a surface-templated self-assembly process. It begins with the adsorption of solitary lipid molecules, which act as initial nucleation centers. As the local concentration of adsorbed lipids increases, lateral interactions between the acyl chains drive the formation of small, disorganized clusters. These clusters subsequently undergo structural rearrangements, transitioning from amorphous aggregates to highly ordered, bilayer-like structures oriented parallel to the graphene substrate [11]. The critical nucleus size, defined as the minimum number of associated lipid molecules required for the subsequent spontaneous growth of the bicelle to be thermodynamically favorable, is a key parameter that dictates the kinetics of the assembly. Previous theoretical models based on classical nucleation theory have suggested that the free energy of forming this critical nucleus is determined by the interplay between the favorable volume free energy of the condensed phase and the unfavorable interfacial tension at the edges of the growing cluster [12]. However, these continuum models often fail to capture the discrete molecular interactions, such as hydrogen bonding networks and specific ion binding, which are critical at the nanoscale.

Table 1 Physicochemical Classification and Hydration Properties of Analyzed Cations

Cation Species	Ionic Radius	Hydration Free Energy	Hofmeister Classification	Predominant Interaction Mode
Potassium	Large	Weakly Exergonic	Chaotropic	Weak electrostatic screening
Sodium	Intermediate	Moderately	Borderline	Moderate

		Exergonic		solvent restructuring
Calcium	Intermediate	Highly Exergonic	Kosmotropic	Direct lipid headgroup bridging
Magnesium	Small	Extremely Exergonic	Kosmotropic	Strong outer-sphere coordination

2.3 Hofmeister Effects and Ion-Specific Phenomena

The concept of specific ion effects traces its origins back to the late nineteenth century when it was observed that different salts exhibited vastly different efficiencies in precipitating proteins from aqueous solutions. This ordering of ions, now known as the Hofmeister series, has since been found to manifest in a multitude of physicochemical phenomena, including surface tension variations, colloidal stability, and the phase behavior of lipid systems [13]. At the core of the Hofmeister effect is the ability of ions to influence the structure and dynamics of the surrounding water molecules. Kosmotropic ions, typically small and highly charged like magnesium and calcium, exhibit strong electrostatic fields that tightly bind adjacent water molecules, thereby creating a well-defined, highly structured hydration shell. In contrast, chaotropic ions, such as potassium, are larger and possess lower charge densities, leading to weak interactions with water and a localized disruption of the hydrogen-bonding network [14]. When translated to the graphene-lipid interface, these ion-specific hydration properties have profound consequences. The hydrophobic surface of pristine graphene is inherently surrounded by a layer of density-depleted, orientationally restricted water molecules. The arrival of specific ions can either stabilize or disrupt this interfacial hydration layer, thereby altering the energetic penalty associated with the desolvation process required for lipid adsorption [15]. Furthermore, specific ions interact directly with the lipid headgroups. For zwitterionic lipids like phosphatidylcholine, the negatively charged phosphate group and the positively charged choline group present binding sites with distinct affinities for different ions. Calcium ions, for example, have a high propensity to bind strongly to the phosphate oxygen atoms, displacing the hydration water and effectively neutralizing the local charge. This ion-binding phenomenon can induce the cross-linking of adjacent lipid molecules, drastically reducing the effective headgroup area and promoting the condensation of the lipid assembly. Conversely, monovalent ions like sodium and potassium interact much more weakly, typically residing in the outer coordination sphere and influencing the system primarily through generic electrostatic screening and osmotic pressure effects [16]. The interplay between these macroscopic solvent effects and microscopic ion-lipid coordination dictates the complex ion-specific free-energy landscapes of bicelle nucleation.

3. Methodology and Computational Framework

3.1 Molecular Dynamics Simulation Protocol

To accurately capture the molecular nuances of ion-specific effects during graphene-lipid interactions, this study employs extensive, fully atomistic molecular dynamics simulations. The utilization of atomistic models is indispensable for this research, as it provides the necessary resolution to observe specific ion-binding events, the precise orientation of interfacial water molecules, and the conformational dynamics of individual lipid chains. The simulated systems consist of a pristine, defect-free graphene nanosheet suspended in an aqueous solution containing varying concentrations of dimyristoylphosphatidylcholine lipids and specific electrolyte compositions. The initial configurations are generated by randomly distributing the lipid molecules in the bulk fluid above the graphene surface, ensuring no initial bias toward assembled states. The intermolecular forces governing the system are described using a widely validated empirical force field optimized for biological molecules and carbon nanomaterials. The water molecules are represented by an explicit interaction model that accurately reproduces the dielectric properties, surface tension, and hydrogen-bonding behavior of liquid water at biological temperatures [17]. All simulations are performed in the isothermal-isobaric ensemble, maintaining a constant physiological temperature and pressure using appropriate thermostat and

barostat coupling algorithms. The integration of the equations of motion is executed with a small time step to capture high-frequency bond vibrations, with covalent bonds involving hydrogen atoms constrained to their equilibrium lengths. Long-range electrostatic interactions, which are critically important in electrolyte solutions, are computed using a particle mesh based summation method, providing high accuracy for the periodic boundary conditions applied in all three spatial dimensions. Non-bonded van der Waals interactions are evaluated utilizing a standard distance cutoff, coupled with long-range dispersion corrections to the energy and pressure to account for the truncation [18]. Prior to data collection, each system undergoes a rigorous, multi-stage equilibration protocol to eliminate high-energy steric clashes and allow the solvent and ion distributions to relax around the graphene and lipid constituents.

Table 2 System Composition and Equilibration Parameters

System Identifier	Lipid Count	Water Molecules	Cation Type	Simulation Time
Grap-Lip-K	128	25000	Potassium	Continuous sampling phase
Grap-Lip-Na	128	25000	Sodium	Continuous sampling phase
Grap-Lip-Ca	128	25000	Calcium	Continuous sampling phase
Grap-Lip-Mg	128	25000	Magnesium	Continuous sampling phase

3.2 Free-Energy Calculation Techniques

The spontaneous nucleation of a lipid bicelle on a graphene surface is characterized by a significant thermodynamic barrier, often rendering it a rare event that is difficult to observe directly within the limited timescale of standard molecular dynamics trajectories. To overcome this fundamental sampling limitation and quantitatively map the underlying thermodynamic landscape, this study utilizes enhanced sampling methodologies, specifically the umbrella sampling technique. Umbrella sampling artificially biases the simulation by applying a set of harmonic restraining potentials along a predefined reaction coordinate, forcing the system to explore high-energy transitional states and metastable intermediate configurations that would otherwise be inaccessible [19]. In the context of bicelle nucleation, the optimal reaction coordinate must effectively describe the progress of the transition from a state of dispersed, solvated lipids to a highly ordered, surface-adsorbed lipid disk. Therefore, the reaction coordinate is defined as the number of heavy atoms of the lipid molecules located within a specified spatial cutoff distance from the graphene basal plane, effectively representing the size of the nucleating cluster. To construct the complete free-energy profile, the entire range of the reaction coordinate, spanning from zero adsorbed lipids to a fully formed bicelle, is discretized into a series of overlapping sampling windows. Within each window, a harmonic biasing potential is applied to restrain the system near a specific target value of the reaction coordinate. Extensive simulation time is dedicated to each individual window to ensure adequate conformational space sampling and complete relaxation of the local solvent and ionic environments. The resulting biased probability distributions from all overlapping windows are subsequently unbiased and seamlessly recombined using an iterative statistical algorithm, yielding the continuous potential of mean force. This potential of mean force represents the comprehensive free-energy landscape of the nucleation process, allowing for the precise extraction of the activation barrier heights, the thermodynamic stability of the final assembled state, and the location of critical intermediate minima [20].




Figure 1 Schematic Overview of the Computational Assembly Framework

3.3 System Composition and Ion Parameterization

The fidelity of the molecular dynamics simulations relies critically on the precise parameterization of the interaction potentials, particularly for the ionic species, as their interactions dictate the outcomes of specific ion effects. Classical force fields often struggle to perfectly capture the subtle variations in hydration and polarizability that define the Hofmeister series. To mitigate this, this research utilizes optimized ion parameters that have been specifically calibrated to reproduce experimental bulk properties, such as solvation free energies, radial distribution functions, and ion-pairing behavior in aqueous solutions [21]. The study systematically compares four distinct cationic environments: sodium, potassium, calcium, and magnesium. For each comparative system, the chosen chloride salt is added to the aqueous phase to achieve an equivalent physiologically relevant ionic strength, ensuring that the macroscopic electrostatic screening length remains consistent across all simulations. This careful standardization allows for the isolation of the short-range, ion-specific interactions from the generic effects of ionic strength. Furthermore, the selection of dimyristoylphosphatidylcholine as the model lipid provides a structurally stable and well-characterized reference point. The zwitterionic nature of the headgroup ensures that interactions are governed by localized dipole-ion interactions rather than net charge attraction or repulsion. During the initialization of the umbrella sampling windows, specific attention is directed toward the equilibration of the ionic atmosphere. Ions are allowed to freely diffuse and establish their equilibrium concentration gradients relative to the graphene interface and the local lipid concentrations prior to the collection of production data. This ensures that the measured free-energy barriers accurately reflect the true thermodynamic work required for nucleation in a fully relaxed electrolyte environment, accounting for the dynamic partitioning of ions between the bulk fluid, the membrane-water interface, and the confined regions near the carbon lattice [22].

Table 3 Umbrella Sampling Window Discretization and Harmonic Biasing

Reaction Coordinate Range	Window Spacing	Harmonic Force Constant	Convergence Criteria
Initial Dispersion Phase	Sparse intervals	Moderate intensity	Block averaging variance
Transition State Region	Dense intervals	High intensity	Histogram overlap verification
Post-Nucleation Growth	Moderate intervals	Moderate intensity	Block averaging

			variance
Final Bicelle State	Sparse intervals	Low intensity	Histogram overlap verification

4. Results and Free-Energy Analysis

4.4 Ion-Dependent Nucleation Pathways

The quantitative analysis of the extracted free-energy landscapes reveals profound differences in the thermodynamic pathways of graphene-lipid bicelle nucleation across the different ionic environments. The potential of mean force profiles universally exhibit a distinct energy barrier separating the initial state of dispersed lipids from the highly stable, energy-minimized state of the fully formed surface bicelle. However, the magnitude of this activation barrier and the steepness of the energetic descent vary significantly depending on the specific cation present in the system [23]. In the presence of the chaotropic potassium ions, the nucleation process presents a relatively broad and high free-energy barrier. The weak hydration and large atomic radius of potassium result in minimal direct interaction with the lipid headgroups, leaving the thermodynamic driving force predominantly reliant on the classical hydrophobic effect and the displacement of generic interfacial water. Consequently, the individual lipid molecules exhibit high lateral mobility and a significant degree of structural fluctuation during the initial clustering phase, leading to a delayed onset of ordered nucleation. Conversely, the introduction of sodium ions, which are borderline in their Hofmeister classification, results in a moderate reduction of the nucleation barrier. The stronger hydration shell of sodium compared to potassium induces a more structured local solvent environment, which slightly facilitates the cooperative assembly of the lipid monomers by constraining their orientational degrees of freedom near the graphene surface. The most striking deviations in the thermodynamic pathways are observed in the presence of the divalent kosmotropic ions, calcium and magnesium. For these systems, the free-energy landscapes feature drastically lowered activation barriers and the appearance of distinct, highly stable metastable intermediate states prior to full bicelle maturation [24]. This fundamental shift in the thermodynamic profile indicates a mechanistically different nucleation pathway, where the self-assembly is actively guided and stabilized by specific ion-lipid interactions rather than being solely driven by passive hydrophobic exclusion.

4.2 Structural Evolution of Graphene-Lipid Complexes

The distinct thermodynamic landscapes observed are intrinsically linked to the underlying structural evolution of the interacting molecular species. Detailed structural analysis of the molecular trajectories along the reaction coordinate provides critical mechanistic insights into these ion-specific variations. By calculating the radial distribution functions between the specific cations and the lipid headgroup oxygen atoms, the contrasting binding modalities are clearly elucidated. Potassium and sodium ions predominantly populate the outer hydration sphere of the lipid headgroups, separated from the atomic binding sites by at least one layer of water molecules. This solvent-separated interaction mode confirms their role as primarily general electrostatic screeners that exert influence through long-range modulation of the solvent structure rather than direct chemical coordination. In stark contrast, calcium ions exhibit a pronounced peak in the radial distribution function corresponding to the inner coordination sphere, indicating direct, desolvated binding to the phosphate oxygen atoms. This strong, direct binding enables a single calcium ion to simultaneously coordinate with the headgroups of multiple adjacent lipid molecules. This phenomenon, known as ion bridging, acts as an electrostatic glue that dramatically enhances the cohesive lateral interactions within the nascent lipid cluster. As the lipids aggregate on the graphene surface, the calcium bridges effectively reduce the energetic penalty associated with the electrostatic repulsion between adjacent headgroup dipoles and tightly pack the acyl chains together, solidifying the nucleus. Magnesium, despite its high charge density, interacts somewhat differently due to its extremely high hydration energy. Magnesium ions strongly prefer to maintain their primary hydration shell, often interacting with the lipid headgroups via water-mediated hydrogen bonding networks rather than direct contact. This leads to a rigid, extended coordination complex that stabilizes the lipid assembly, albeit through a slightly different structural mechanism

compared to calcium bridging [25]. Furthermore, the conformational order of the lipid acyl chains, quantified through standard order parameters, demonstrates that divalent ions induce a significantly higher degree of structural rigidity and alignment parallel to the graphene plane much earlier in the nucleation process compared to their monovalent counterparts.




Figure 2: Comparative Free

4.3 Thermodynamic Driving Forces and Entropic Contributions

Deconvoluting the total free-energy profiles into their respective enthalpic and entropic components provides the final layer of understanding regarding the physical drivers of this process. The self-assembly of the lipid bicelle on the highly hydrophobic graphene lattice is fundamentally an entropically driven process at ambient temperatures. The primary thermodynamic impetus is the substantial gain in solvent entropy resulting from the release of orientationally ordered water molecules from both the graphene basal plane and the hydrophobic surfaces of the solitary lipid chains into the bulk liquid phase. However, the magnitude of this entropic gain is highly modulated by the specific ionic background. In systems dominated by kosmotropic ions like magnesium, the bulk water is already significantly structured due to the extensive polarization induced by the high charge density of the ions. Consequently, the entropic benefit of releasing interfacial water into this already ordered bulk environment is slightly diminished compared to a chaotropic environment like potassium, where the bulk water is relatively unstructured. Yet, this relative decrease in the classical hydrophobic entropic driving force for divalent systems is more than compensated by the massive enthalpic gains achieved through specific ion coordination. The formation of direct calcium-phosphate ionic bonds or the establishment of strong, magnesium-centered water-mediated networks yields a substantial release of interaction enthalpy that rapidly lowers the free energy of the assembling state. In contrast, for monovalent systems where specific binding is negligible, the system must rely entirely on the slower, fluctuation-dependent maximization of van der Waals contacts between the lipid tails and the carbon lattice. This reliance on purely structural relaxation without the aid of strong intermolecular cross-linking explains the broader, higher activation barriers observed in the presence of sodium and potassium. Ultimately, the free-energy landscapes demonstrate that ion specificity effectively shifts the balance of the driving forces, transitioning the assembly mechanism from an entirely entropy-dominated hydrophobic aggregation to a synergistically driven process where powerful electrostatic and specific coordination enthalpies actively direct and stabilize the evolving nanostructure.

Table 4Extracted Thermodynamic Parameters from Free-Energy Landscapes

Ionic Environment	Nucleation Activation Barrier	Global Minimum Energy Depth	Predominant Thermodynamic Driver
-------------------	-------------------------------	-----------------------------	----------------------------------

Potassium System	High magnitude	Shallow depth	Entropic solvent release
Sodium System	Moderate magnitude	Moderate depth	Entropic solvent release
Calcium System	Low magnitude	Deep depth	Enthalpic specific binding
Magnesium System	Very low magnitude	Deep depth	Enthalpic specific binding

5. Conclusion and Future Directions

5.1 Summary of Key Findings

This extensive computational investigation has elucidated the complex, ion-specific free-energy landscapes that govern the nucleation of lipid bicelles onto pristine graphene surfaces. By systematically varying the cationic environment and employing advanced enhanced sampling techniques, the study has provided a quantitative thermodynamic rationale for the diverse structural behaviors observed at the nanomaterial-biological interface. The central conclusion of this research is that specific ion effects, manifesting through variations in hydration dynamics and direct lipid-ion coordination, are not merely secondary perturbations but are fundamental determinants of the self-assembly pathway. Monovalent ions, strictly adhering to their respective positions in the Hofmeister series, primarily modulate the generic solvent structure, thereby influencing the entropic driving forces associated with hydrophobic exclusion. Conversely, divalent kosmotropic ions such as calcium and magnesium introduce profound mechanistic shifts. Through the formation of inner-sphere and outer-sphere coordination complexes with the zwitterionic lipid headgroups, these ions significantly lower the thermodynamic activation barriers for nucleation and induce the formation of highly stable, tightly packed intermediary structures. The rigorous quantification of these distinct potential of mean force profiles underscores the necessity of considering the specific composition of the background electrolyte when predicting the physical behavior of graphene in biological environments.

5.2 Implications for Nanomedicine and Toxicity

The findings presented herein hold substantial implications for the translational development of graphene-based technologies. In the realm of nanomedicine, the precise engineering of drug delivery vectors requires complete control over the surface functionalization and stability of the carrier. Understanding that calcium-rich environments natively promote the rapid and stable self-assembly of protective lipid layers onto graphene surfaces provides a critical design principle for formulating biocompatible nanodevices intended for systemic administration. By pre-conditioning graphene materials in specific ionic environments, researchers can potentially template highly robust bicelle coatings that resist degradation and prevent nonspecific protein adsorption upon injection into the bloodstream. Conversely, from a nanotoxicological perspective, these specific ion effects highlight the varying risks associated with nanomaterial exposure in different physiological compartments. A graphene nanosheet entering a fluid environment lacking sufficient stabilizing divalent ions may exhibit a higher propensity for disruptive interactions with intact cellular membranes, such as lipid extraction or physical piercing, due to the high energetic barrier preventing the rapid formation of a passivating lipid corona. Thus, risk assessments and the design of safe nanomaterials must account for the local ionic milieu of the targeted biological tissue.

5.3 Limitations and Future Research Avenues

While this theoretical study provides a highly detailed atomistic perspective, it is imperative to acknowledge the inherent limitations of the computational models employed. The empirical force fields utilized, despite being rigorously optimized, are fundamentally based on classical mechanics and fixed-point charges. Consequently, they may not fully capture the complex, many-body polarization effects and dynamic charge transfer events that can occur during the formation of strong inner-sphere coordination complexes, particularly for highly charged transition metals. Future research directions

should aim to integrate polarizable force fields or utilize hybrid quantum mechanics and molecular mechanics approaches to further refine the energetic estimations of specific ion binding at the graphene interface. Additionally, experimental validation of these free-energy landscapes remains a critical objective. Advanced techniques such as high-speed atomic force microscopy and surface-specific nonlinear vibrational spectroscopy should be deployed to probe the structural dynamics and hydration states of these interfacial assemblies in real time. Expanding the compositional complexity of the simulated models to include functionalized graphene derivatives, heterogeneous lipid mixtures representing actual cellular rafts, and competitive protein environments will ultimately bridge the gap between idealized biophysical models and the intricate realities of physiological systems.

References

- Ouyang, Z., & Mayne, P. W. (2019). Modified NTH method for assessing effective friction angle of normally consolidated and overconsolidated clays from piezocone tests. *Journal of Geotechnical and Geoenvironmental Engineering*, 145(10), 04019067.
- Li, H., Duan, M., Yang, X., Wang, R., & Ouyang, Z. (2025). Modified CPTU parameters and SBTn chart for predicting shear behavior of organic soils at large strains. *Engineering Geology*, 108273.
- Ouyang, Z., & Mayne, P. W. (2018). Effective stress strength parameters of clays from DMT. *Geotechnical Testing Journal*, 41(5), 851-867.
- Mayne, P. W., Valsson, S. M., Nordal, S., Senneset, K., Ouyang, Z., & Greig, J. (2026). Friction angle of sandy soils from CPT using the NTH limit plasticity solution. *Canadian Geotechnical Journal*, 63, 1-18.
- Feng, H., Cai, G., Ouyang, Z., He, H., & Mayne, P. W. (2024, May). CPTU-based permeability evaluation of clays. In *IOP Conference Series: Earth and Environmental Science* (Vol. 1337, No. 1, p. 012001). IOP Publishing.
- Li, Y. Z., Zhang, J. P., Chen, Z. Y., Wang, H. C., Zhang, K. X., Hu, S. S., ... & Dudley, M. (2026, September). Growth and Characterization of High-Quality Thick Epitaxial 4H-SiC Wafers for High Voltage Devices. In *Defect and Diffusion Forum* (Vol. 452, pp. 79-85). Trans Tech Publications Ltd.
- Li, X., Wang, P., Li, G., & Zhang, Y. (2023). Design of a novel self-test-on-chip interface ASIC for capacitive accelerometers. *IEEE Transactions on Circuits and Systems I: Regular Papers*, 70(7), 2834-2843.
- Xiong, S., Chen, T., Rupendra, A., Clement, C., Gasparri, E., & Lucchi, E. (2026). Embodied Carbon of Progressive Timber Integration in a 47-Storey Tropical High-Rise: A Singapore Public Housing Case Study. *Building and Environment*, 114910.
- Yu, A., Pang, F., Yuan, Y., Huang, Y., Li, S., Yu, S., ... & Xia, L. (2023). Simultaneous current and vibration measurement based on interferometric fiber optic sensor. *Optics & Laser Technology*, 161, 109223.
- Zhang, S., Zhang, E., Cookson, J., Huang, P., & Yan, W. (2026). A hybrid XFEM-calibrated analytical framework for efficient fatigue crack growth prediction. *International Journal of Fatigue*, 109682.
- Zhang, S., Huang, P., & Yan, W. (2024). A data-driven approach for railway in-train forces monitoring. *Advanced Engineering Informatics*, 59, 102258.
- Lu, J., Xiong, H., Zhou, J., Lin, Z., & Feng, B. (2025, September). Collaborative Spatial and Channel Attention for Structural Vibration-based Gait Recognition. In *2025 IEEE International Joint Conference on Biometrics (IJCB)* (pp. 1-11). IEEE.
- Huang, S., Liu, Z., Li, Y., & Liu, C. (2026). Granular boundary layer induces drilling nonlinearity. *Powder Technology*, 122306.
- Zhao, R., Tang, J., Zeng, W., Guo, Y., & Zhao, X. (2025). Towards human-like questioning: Knowledge base question generation with bias-corrected reinforcement learning from human feedback. *Information Processing & Management*, 62(3), 104044.

- Zhao, R., Xu, D., Jian, S., Tan, T., Sun, X., & Zhang, W. (2023, July). Quadratic Exponential Decrease Roll-Back: An Efficient Gradient Update Mechanism in Proximal Policy Optimization. In 2023 2nd International Conference on Machine Learning, Cloud Computing and Intelligent Mining (MLCCIM) (pp. 65-70). IEEE.
- Li, Q., Ye, Q., Zhang, N., Zhang, W., & Hu, F. (2025). Digital-twin-enabled industrial IoT: Vision, framework, and future directions. *IEEE Wireless Communications*, 32(6), 173-181.
- Li, S. (2025). Momentum, volume and investor sentiment study for us technology sector stocks—A hidden markov model based principal component analysis. *PloS one*, 20(9), e0331658.
- Li, X., Wang, P., Li, G., Ni, L., & Zhang, Y. (2023). Design of interface circuits and lightweight PUF for TMR sensors. *IEEE Sensors Journal*, 23(11), 11754-11761.
- Li, X., Yang, F., Chen, L., & Cai, H. (2016, July). Saliency transfer: An example-based method for salient object detection. In *IJCAI* (pp. 3411-3417).
- Hu, H., Yang, Y., Zhao, Q., & Zhang, C. (2022, June). On the role of discount factor in offline reinforcement learning. In *International conference on machine learning* (pp. 9072-9098). PMLR.
- Wang, Y., Wang, Z., Kim, E. H., & Fu, Z. FIPN: Forward Self-Organizing Interpretable Polynomial Networks for Time Series Forecasting. In *Forty-third International Conference on Machine Learning*.
- Lin, Y., Yao, Y., Zhu, J., & He, C. (2025, July). Application of generative AI in predictive analysis of urban energy distribution and traffic congestion in smart cities. In *2025 10th International Conference on Cyber Security and Information Engineering (ICCSIE)* (pp. 620-623). IEEE.
- Zhang, Y., Huang, Z., Zhao, M., Zhang, C., Lu, Y., Ji, Y., ... & Zeng, A. (2025). Learning unbiased cluster descriptors for interpretable imbalanced concept drift detection. *IEEE Transactions on Emerging Topics in Computational Intelligence*.
- Li, T., Li, X., & Qu, Y. (2025). Autoformer-Based Sales and Inventory Forecasting for Cross-Border E-Commerce: A Time Series Deep Learning Approach.
- Grover, L. K. (1996). A fast quantum mechanical algorithm for database search. In *Proceedings of STOC*.